

SB 608
.S95 T3
Copy 1

The Diseases of the Sweet Pea

THESIS

PRESENTED TO THE FACULTY OF THE GRADUATE SCHOOL OF
THE UNIVERSITY OF PENNSYLVANIA IN PARTIAL FUL-
FILMENT OF THE REQUIREMENTS FOR THE
DEGREE OF DOCTOR OF PHILOSOPHY

J. J. TAUBENHAUS

PHILADELPHIA

1914



The Diseases of the Sweet Pea

THESIS

PRESENTED TO THE FACULTY OF THE GRADUATE SCHOOL OF
THE UNIVERSITY OF PENNSYLVANIA IN PARTIAL FUL-
FILMENT OF THE REQUIREMENTS FOR THE
DEGREE OF DOCTOR OF PHILOSOPHY

acoh
J. J. TAUBENHAUS



PHILADELPHIA

1914

SB 608
.S95T3

The University
of Chicago



BULLETIN NO. 106

NOVEMBER, 1914

Delaware College
Agricultural
Experiment Station



The Diseases of the Sweet Pea

BY J. J. TAUBENHAUS

Newark, Delaware

INDEX

The Host	3—10
Taxonomy and Botany	3—
History of the Sweet Pea	5— 6
Economic Importance	5— 6
Cultivation and Care	6— 8
Sweet Peas Under Glass	8—10
Diseases of the Sweet Pea	10—74
I. Fungous Diseases	12—68
II. Bacterial Diseases	68—72
III. Physiological Diseases	72—74
A. Root Diseases	12—36
B. Diseases of the Aerial Parts of the Plants.....	36—65
C. Diseased Seeds	65—68
Root rot (<i>Thielavia basicola</i> Zopf)	12—21
Rhizoctonia root rot (<i>Corticium vagum</i> B. & C.)	21—27
Chaetomium root rot (<i>Chaetomium spirochaete</i> Patt.)..	27—30
Fusarium root rot (<i>Fusarium lathyri</i> Taub.).....	31—34
Root Knot (<i>Heterodera radicolica</i> [Greef] Muller).....	35—36
Stem or collar rot (<i>S. libertiana</i> Fekl.)	36—41
Powdery mildew (<i>Oidium</i> sp.)	41—43
Anthracnose (<i>Glomerella rufomaculans</i>)	43—55
Mosaic disease	56—64
Diseases not known to be present in this country.....	64—65
Pea blight (<i>Peronospora trifoliorum</i> De By.)	64—65
Pea spot (<i>Ascochyta pisi</i> Lib.)	65
Diseased seeds	65—68
Streak disease (<i>Bacillus lathyri</i> M. & T.)	68—72
Bud drop	72—73
Arrested development	73—74
Methods of Control	74—76
Resistant varieties	76—80
Seed treatment	76—80
Treatment of Soil with Chemicals	80—82
Studies of the Fungicidal value of some chemical poisons	82—85
Soil treatment in the greenhouse	85—87
Summary	85—89
References	88—93

THE DISEASES OF THE SWEET PEA*

BY J. J. TAUBENHAUS

THE HOST

TAXONOMY AND BOTANY

The sweet pea (*Lathyrus odoratus* L.) belongs to the family Leguminosae, the sub family Papilionaceae, the tribe Viciae. The sweet pea (*Lathyrus odoratus* L.) is a herbaceous vine with rough stems, hairy and winged. The leaves are alternate, pinnately compound with terminal tendriform leaflets. The leaflets are oval or oblong, mucronate. Peduncles 2-4 flowered, much longer than the leaves. The calyx teeth are broad, longer than the tube. The flowers are large, and showy in shades of blue, red, yellow and white. The standard is large, expanded, hooded, or wavy. The legumes are compressed, linear, 1-3 inches, hairy. The seeds are round, sometimes angled, black, white or mottled.

HISTORY OF THE SWEET PEA

*Origin, Improvement and Distribution*₁**.

The word "Lathyrus" is from the Greek *La. la* (augmentative) and *thouros*, anything exciting, having reference to the qualities of seeds of certain species. In Europe the species of *Lathyrus* are known as "Gesse," the sweet pea being known as *Gesse odorante*. The French know the plant under this name, or as *Pois Odorante*, or *Pois de Senteur*."

The earliest mention of the sweet pea is found in "Syllabus Plantarum Sicillæ-nuper detectarum a P. F. Franciscus Cupani" (Panormi, 1695). The sweet pea is spoken of as "*Lathyrus distoplatyphyllos hirsutis mollis, magno et peramoeno flore odore*." Father

*Also presented to the Faculty of the Graduate School of the University of Pennsylvania, June 1913, as a major thesis in partial fulfillment of the requirements for the degree of Doctor of Philosophy.

Acknowledgements.—The writer is indebted to Dr. John W. Harshberger, under whose direction the work was conducted, for helpful suggestions and criticisms. Acknowledgements are also due to Dr. T. F. Manns for helpful suggestions and advice. Thanks are also due to the many American seedsmen for financial support in carrying out the field experiments.

**All references will be found on pp. 84 to 93.

Cupani was very enthusiastic about this flower and in 1699 sent seed to Dr. Uvedale at Enfield, England, and to Caspar Commelin at Amsterdam, Holland. Commelin described and illustrated the plant in his "Hort.-Medici Amstelodamensis" (1697-1701). Commelin also adopted Cupani's name for the plant.

In his "Almagesti Botanici Mantissa" (1700) Dr. Leonard Plukenet also gives a description of the sweet pea. A dried specimen preserved in Plukenet's Herbarium, which now forms part of Sir Hans Sloane collection, is the oldest specimen of the sweet pea in existence.

Mention is made of the sweet pea by Petiver in the "Botanicum Hortense III" (1713). Petiver calls the plant *Lathyrus Siculus*, a native of Sicily which has large broad sweet smelling flowers. H. B. Ruppil in "Flora Jenensis" (Frankfort, 1718) places *Lathyrus Siculus Ravini* in a class of plants with irregular flowers. It is thus seen that all authorities place Sicily as the home of the sweet pea.

Linnaeus, 1753, in his great "Systema Plantarum Europæ," classifies the sweet pea as follows:

"Odoratus II. *Lathyrus pedunculis bifloris, cirrhis diphyllis, foliis ovato-oblongis, leguminibus hirsutis.* Hort. Cliff. 368, Hort. Upsal. 216, Roy. lugd. 363.

"*Siculus a. Lathyrus Siculus.* Rupp. jen., 210

Lathyrus distoplatyphyllos hirsutis mollis, magno et peramæno flore odoro. Comm. Hort. 2, p. 219, t. 80.

"*Zeylanicus b. Lathyrus Zeylanicus. Odorato flore amœne ex albo et rubro vario.* Burm. Zeyl., 138.

"Habitat: a. in Sicilia; b. in Zeylona."

Here then is the first use of the term "odoratus" as a distinctive name.

Kniphof in his "Botanico in originali" (1757-1763) gives a colored illustration of Painted Lady sweet pea. In the catalog of W. Malcolm (1778), seedsman of Kensington Turnpike, we find offered for sale, white, purple and Painted Lady sweet peas. The first evidence of improvement is noticed in the catalog of John Mason (1793). He offered black, purple, scarlet, white and Painted Lady sweet peas. Between 1845 and 1849 the firm of Messrs. J. Carter & Co. introduced a new striped sweet pea and a new large purple sweet pea. In 1850 Messrs. Nobel, Cooper and Bolton introduced a new large dark purple variety. In 1860 Mr. Carter offered several new varieties of sweet

peas. In James Vick's "Illustrated Catalog and Flower Guide" (1870), nine varieties of sweet peas are mentioned.

Beginning with 1880, great strides have been made in the improvement of the sweet pea in England. Thomas Laxton and Henry Eckford (about 1880) were the moving spirits. Mr. Laxton introduced several new varieties obtained by crossing. Mr. Eckford was responsible for one hundred and fifteen new varieties. In America, Edward Sayers in his book "The American Flower Garden Companion" Boston, 1838), gives a list of five varieties of sweet peas. Of the American pioneers and breeders of the sweet pea, those who should be mentioned are D. M. Ferry & Co. who in 1889 introduced the Blanche Ferry; W. Atlee Burpee & Co., Messrs. C. C. Morse & Co., J. C. Vaughn and Peter Henderson.

During the first one hundred years of sweet pea culture only three varieties, or colors, were known, i. e., purple with blue wings, pale red with white wings (Painted Lady) and white. The black and the scarlet appeared in the last years of the eighteenth century. At the present time there are more than a hundred and fifty varieties in cultivation with promising new ones appearing every year. This shows the great popularity of the sweet pea and the extent to which it is grown. Whenever a crop is grown extensively and for a long time under certain soil and climatic conditions, as is the sweet pea, diseases are sure to appear, making it difficult for the crop to succeed unless precautionary measures are taken. The cultivation of the sweet pea in England is at a crisis, the disease factor being the one obstacle to its cultivation. In America sweet pea growers are confronted with several important diseases.

ECONOMIC IMPORTANCE

I have been unable to obtain statistical data concerning the sweet pea crop. Messrs. C. C. Morse & Co. furnished us with the following information:

"Your favor of the 8th inst. was duly received and I shall be very glad to answer the questions you have asked, as well as I can.

"However, our Sweet Peas will be practically a failure this year and my statistics will apply only to past seasons, probably more accurately to the crop of 1911, which was the best we have had in recent times. Even last year's crop was very poor.

1. So far as seed is concerned, the Sweet Pea crop is worth about \$250,000 annually to the grower.

2. There are about 1700 acres of land planted to Sweet Peas for seed in California annually. I have no knowledge of what acreage is devoted to flowers for market.

3. Practically no Sweet Pea seed is imported and about one-half of the California acreage is exported.

4. Fully 90 per cent. of the export business is done with Great Britain; the balance with Holland, Germany and France.

5. No other country, so far as we know, produces Sweet Pea seed to amount to anything.

Respectfully yours,

C. C. MORSE & CO."

According to Bailey², California, in 1902, supplied the world's market with 125 tons of sweet pea seeds. As a cut flower the sweet pea is a great favorite and is extensively grown for that purpose.

CULTIVATION AND CARE

In this connection we will consider only those points of culture which directly influence the disease factor.

Climate. The sweet pea does best in a temperate region. It will not stand too warm a climate, as the plants there soon dry up and die, or they are so weakened as to succumb readily to all sorts of fungous diseases. California seems to be its ideal home, nevertheless the sweet pea is known to thrive under various climatic conditions. It is less susceptible to cold than to heat and in hot dry climates irrigation is essential.

Site. The sweet pea requires an open, sunny location so as to get plenty of light and air. Plants grown in too shady a place will be spindly, weak, and open to the attacks of diseases.

Soil. All light sandy soils should be avoided for the reason already referred to above. A good loamy soil is preferred provided, of course, its subsoil is well underdrained, otherwise the plants will grow poorly and be constantly open to the attacks of disease.

Fertilizer. In order to be at their best, sweet peas must be provided with sufficient available plant food in the soil. However,

fertilizers should be used very judiciously. The aim should be to apply a food that is well balanced, i. e., it should contain the proper amount of nitrogen, phosphorus, potash and lime. Too much of one of these elements and too little of the other will produce disturbances in the metabolism of the plant as will be seen later under the discussion of physiological diseases. For an ordinary garden land, the following is a well balanced fertilizer devised by Prof. T. F. Manns of the Del. Expt. Station. Before plowing a surface application of well rotted manure at the rate of 6 tons per acre is first applied to the soil. After plowing and harrowing the soil, it is furrowed and 5 tons of rotted manure per acre is applied in the furrows. The manure is worked in deep with a spade and the following fertilizer is applied in the furrows:

*Sodium nitrate	200 lbs. per acre
Dried blood	250 lbs. " "
Acid phosphate	1200 lbs. " "
Potassium sulphate	240 lbs. " "
Rock phosphate	400 lbs. " "
Hydrated lime	200 lbs. " "
Carbonate of lime	600 lbs. " "

The fertilizer is well mixed up with the soil and the seeds are planted on top and covered to a depth of about two to three inches.

Care of the Seeds and Depth of Sowing. Most of the white-seeded varieties are subject to decay in the soil. Most of the black-seeded varieties are more resistant to soil decay but they do not germinate evenly. In order, therefore, to hasten germination, it is advisable to place the seed in tepid water over night. With this treatment the seeds swell and are ready to be sown the next day. The Rev. W. T. Hutchins, a well-known sweet peas specialist in America, advises the placing of the seed in moist earth for seven or eight days. They are then taken out and examined. The swollen seeds are planted and the hard seeds cut with a knife to hasten germination. Whatever method is used the aim should be to hasten germination in order to prevent the seed from laying too long in the ground and thereby causing decay. The depth of sowing the seeds varies from two to three inches according to the nature of the soil. As to distance, five feet

*This mixture was recommended for a very heavy acid soil deficient in organic matter.

apart between the rows and three inches in the row will insure the desired amount of air and light.

Care of the Growing Plants. Frequent cultivations with the hoe or with the cultivator will provide sufficient aeration of the roots to insure a vigorous growth of the vine. It is in baked and ill-drained soils that saprophytic fungi assume the nature of semi parasites, since it is in these soils that the plants are often weak and consequently yield readily to the attacks of disease. Moreover, frequent cultivations destroy the weeds which may act as disease carriers or disease trans-



A convenient way of trailing sweet peas

mitters. Irrigation wherever possible will no doubt benefit the plants, but irrigation should not replace cultivation. The plants should be kept free from the insect and the fungus pests. This will be discussed under methods of control. With the sweet pea, contrary to many other flowering plants, the blooms should be gathered freely as the more we do this the longer the vines will continue to flower.

SWEET PEAS UNDER GLASS

The following notes on cultivation are by Mr. William Sim, Clifftondale, Mass., and are extracted from a paper read by him before the Gardeners' and Florists' Club of Boston on April 21, 1908:

“To grow the sweet pea to perfection under glass you must have a greenhouse suitable for the purpose. It should be at least eight feet high on the sides, four and a half feet being glass. My houses are seven feet, and I find the side rows strike the glass when the vines are about half grown, thereby giving me half a crop. My center rows are about right; they are twelve to fifteen feet high. The higher the vines grow the more and better flowers you get. We plant the rows five feet apart and in a line with the supports of the greenhouse. The up-rights are twelve feet apart, so in supporting we run twine from one support to the other on each side of the row. This I have found the best method of supporting. I have tried wire netting; it is only a nuisance, as the vines do not cling to the wire, which causes just as much tying as if it were not there. It also causes injury many times to the vines, as a sweet pea stretches many times more than a foot in developing; if held back by anything in growing the growth looks like a spiral spring, and the picking of the blooms is made very difficult. The side rows are planted five feet from the sides of the house; and all the heating pipes are on the sides. The vines are very susceptible to red spider and as they will not stand syringing, the further you can afford economically to have them from the pipes the better.

“We have not changed the soil in the houses since they were built four and five years ago, and we find the vines are getting more vigorous each year. In the same soil a crop of tomatoes and of violets is harvested each year. The soil was originally eighteen inches deep, but by the application of manure each year the depth is now two and a half feet. The tomato crop is on the wane the middle of August. When these are cleaned out we trench the house over as deep as the soil, bringing the bottom soil to the surface. In the bottom of the trench we put three inches of decomposed cow manure; one foot from the surface we put on three inches more of the same material. The house is allowed to remain in this state until nearly time for sowing the seed. The soil is then usually very dry, so we dampen it down enough to cling together while the house gets another forking over. This time we go down one foot and mix the top layer of manure with the surface soil. We then make the surface as nearly level as pos-

sible and thoroughly water the soil, giving enough to penetrate the entire mass, with a strong dose of liquid horse manure. In about three days, depending on the weather, the house will be ready to plant. We sow the seeds about one and a half inches apart. We make the drills one inch deep and do not allow more than one inch of soil over them. We do not pull any more soil toward the roots, as is often recommended, but let it remain level. If more soil is pulled around the base of the plant, stem rot is sure to follow. We do not water the plants again until they are up about three inches.

“Of course, you can grow them on a bench with a few inches of soil, but the results will be just what you make them—a weak growth and a crop of short-stemmed flowers. These soon play out, as there is not enough soil or food for the vines to live on.

“They may be made to flower any time you wish by increasing the temperature, but the best results are obtained by growing at a temperature just above freezing until the buds can be felt in the crowns of the plants. Then the temperature should be gradually increased, say one degree a night, until you reach 48 degrees. This, I think, is about right, although in midwinter I think they move a little better at 50. As the days lengthen a little cooler temperature seems to suit better. A rise of 10 to 15 degrees should be given during the day in sunny weather. In spells of cloudy weather 55 degrees is high enough during the day. If a high temperature is given in dark weather the growth gets soft and wilts when the sun comes out bright again. While the plants are young they should be regularly fumigated so there will not be a sign of lice when the plants commence to flower. If they are clean at this stage it will not be necessary to fumigate while they are in bloom. It is impossible to sell sweet peas that smell of tobacco. Tobacco also bleaches the flowers of some varieties, and makes them look like some other variety.

“We sometimes hear of someone having trouble with the buds dropping. This is more the case in midwinter than at any other time, and is caused by a too cool temperature or a sudden chill, or too much water. Should a house be allowed to go near the freezing point in midwinter the wholesale dropping of buds will be sure to follow.”

DISEASES OF THE SWEET PEA

HISTORICAL

The literature on the subject of sweet pea disease previous to 1906 is mostly of a fragmentary nature. In 1896 Cuthbertson,³ first records a bud and blossom drop of the Cupid sweet pea in Scotland attributing the cause to cold and damp weather at that time.

In 1906, Masee,⁴ gave the first brief scientific account of some sweet pea diseases, mentioning the following fungi: *Peronospora trifoliorum*, *P. viciae*, *Erysiphe polygoni*, and *Ascochyta pisi*.

In 1906, Weston,⁵ was first to describe the "Streak" as a new disease of the sweet pea in England (the cause not given). In 1907, Weston,⁶ again calls attention to the serious nature of the "Streak."

In 1907, an anonymous note,⁶ mentions the following diseases: eelworm (*Tylenchus devastatrix* and *obtusa*), *Peronospora trifoliorum*, sclerotia of some species of *Sclerotinia*, *Erysiphe martii*, and *Botrytis cinerea*.

In 1909, in a brief note, Masee,⁷ also mentions the "Streak" disease which he thinks is induced by an excess of manure in the soil. This excess produces a deleterious effect on the soil flora which in turn brings about physiological disturbances resulting in the "streak." In 1912, Masee,⁸ again mentions a disease of sweet pea seedlings and of other plants as due to *Thielavia*.

In the same year (1912) Chittenden,⁹ before the London Sweet Pea Society and in an article in the Royal Horticultural Society Journal reports on the "Streak" disease, which according to him, was found to be due to *Thielavia basicola*.

In 1912, W. Dyke,¹⁰ an amateur scientist and gardener, calls attention to the "streak" disease which he believes is induced by a species of *Fusarium* and *Macrosporium*. It will be seen from the above reference that the only ones of scientific importance are those of Masee and Chittenden, because both of these investigators base their facts on research. However, as it will be shown later, both Masee and Chittenden mistook *Thielavia* as the cause of the "Streak" disease.

In American literature, Sheldon,¹¹ was the first one to call attention to the anthracnose of the sweet pea (*Glomerella rufomaculans*.)

The diseases of the sweet pea have received no other attention at the hands of American plant pathologists.

For the past three years the writer has been investigating the diseases of the sweet pea, and as a result three papers have already been published_{1,2}.

It is the purpose of the present thesis to bring together all the results obtained in my investigations. The subject is by no means exhausted, as yet, and we hope to devote many more years to the study of the diseases of the sweet pea.

THE DISEASES OF THE SWEET PEA

The diseases treated in this thesis are as follows:

- I. Fungous Diseases.
- II. Bacterial Diseases.
- III. Physiological Diseases.
- IV. Animal or Insect Pest (Of the animal pests I will only consider the *Heterodera radiculicola* and will discuss it under "root diseases." Of the insect pests I will only consider the green aphids and these will be discussed in relation to the "mosaic.")

I. Fungous Diseases

- A. Root Diseases.
- B. Diseases of the Aerial Parts of the Plants.
- C. Diseased Seeds.

A. Root Diseases

All root troubles of the sweet pea are caused by fungi which live primarily in the soil. They can, therefore, also be designated as soil diseases. Diseased roots invariably indicate an infected soil. All soil parasites are not necessarily confined to the roots of the sweet pea only, as we shall have occasion to show later. Of the soil organisms which attack the roots, the following have been investigated:

- Root rot. (*Thielavia basicola* Zopf.)
- Root rot. *Rhizoctonia* (*Corticium Vagum* B. & C.)
- Root rot. (*Chaetomium spirochaete* Patt.)
- Root rot. (*Fusarium lathyri* n. sp.)
- Root galls Eel worms (*Heterodera radiculicola*.)

ROOT ROT (*Thielavia basicola* Zopf)

Historical, Synonymy and Relationship. *Thielavia basicola* belongs to the ascomycetous family Perisporiaceæ. The fungus was

first described by Berkeley and Broome¹³ in 1850, who gave it the name of *Torula basicola*; they found it growing at the base of affected pea plants. The torula or chlamydospore stage is the most conspicuous and it is abundantly found on the host. In 1875, Zopf¹⁴ found the fungus on roots of *Senecio elegans* in Berlin, and in 1876, Sorokin¹⁵ found it on the roots of *Cochlearia armoracia* (horse radish) in Russia, and named it *Helminthosporium fragile*. Zopf, however, made a more thorough study of the fungus, and he discovered the perfect stage, placing it in the Perisporiaceæ, creating a new genus *Thielavia* after Prof. F. von Thielav of the University of Berlin.

In 1886 Saccardo¹⁶ found the same fungus. Noting the observations of Sorokin, he did not agree with him in calling the fungus *Helminthosporium*, and he placed it in the genus *Clasterosporium*. Saccardo thus failed to identify Sorokin's *Helminthosporium fragile* with the chlamydospore stage of *Thielavia* and with Berkeley and Broome's *Torula basicola*. It was to the credit of Sorauer¹⁷ who discovered the relationship of these different stages to belong to one and the same fungus. The name of the fungus with its synonymy is as follows:

- Thielavia basicola* (B. & R.) Zopf.
- Torula basicola* (B. & R.)
- Helminthosporium fragile* Sor.
- Clasterosporium fragile* (Sorok.) Sacc.

Thielavia a Parasite on Other Hosts. The table on page 12 will show the list of hosts parasitized by *Thielavia* together with the authority for the same.

Thielavia Attacking Sweet Peas. In 1912, Chittenden¹⁸ was asked by the National Sweet Pea Society of England to investigate the dreaded "Streak" disease of the sweet pea. In his report before that society Chittenden gives an accurate description of the "Streak," so that there can be no doubt but that he had the disease well in mind, that is, he described it as a stem disease. Chittenden found *Thielavia basicola* to be the cause of "Streak" as he states, "careful microscopic examinations of the brown patches of the roots, and one must insist on the need for care; hasty examination may fail to reveal anything showing that they were attacked by a fungus which turned out to be *Thielavia basicola*. The same fungus was present in practically

<i>Name of Plant</i>	<i>Authority</i>
Aralia quinquefolia	Selby
Begonia rubra	Selby
“ sp.	Zopf
Catalpa speciosa	Selby
Cochlearia armoracia	Sorokin
Cyclamen	Sorokin
Gossypium herbaceum	Smith, E. F.
Lupinus albus	Zopf
Linaria canadensis	Gilbert
Nemaphila auriculata	Berkeley and Broome
Nicotiana tabacum	Selby, Gilbert, Clinton
Onobrychis crista—galli	Zopf
Oxalis corniculata	Gilbert, Stewart
Pisum sativum	Berkeley and others
Phaseolus vulgaris	Zopf
Trigonella coerulea	“
Vigna sinensis	Smith, E. F.
Viola odorata	Thaxter and others
Trifolium repens	Gilbert and Stewart
Lathyrus odoratus	Taubenhaus

every case, sometimes abundantly fruiting, sometimes with only a few spores.”

I will show later where Chittenden has erred in his investigations, and that the “streak” is not a root but a stem disease which is induced by a bacterium and not by Thielavia. In every case where Thielavia has been reported as a parasite on other hosts, it has been found on the roots and not on the stem. This same holds true for the sweet pea. Moreover, Chittenden in his artificial inoculation with the fungus, has succeeded in reproducing the typical root rot and not the “streak” on the stems. Masee¹⁹ too made the same mistake as Chittenden, for he too considers Thielavia to be the cause of the “Streak.” I have seen the Thielavia root rot on half an acre of sweet peas at the trial grounds of one of our commercial seed men. The plants on that infested area were carefully examined, and no signs of streak could be found on the stems. On the other hand, the trouble was seen to be plainly localized at the roots.

Symptoms of Thielavia Root Rot on Sweet Peas. Plants severely infected have practically little or no root system since the latter is destroyed by the fungus as rapidly as the roots appear. (Figs. 1 and 2). Whatever root system is present is of a stubby nature and



Fig. 1*. Root rot caused by *Thielavia*. Contrasting a healthy with a diseased plant of the same age.

* Figs. 1, 2, 9, 17, 33, 34, 35, 37, 39, 40 and 42, Electrotypes, Gardener's Chronicle (England). Photographs by the author.

charred in appearance. The fungus sometimes works upon the stem to a distance of two to three inches above ground, but never to the extent of invading the entire stem. It is probably due to this that some workers have mistaken this disease for the well-known "streak."



Fig. 2. Root rot caused by *Thielavia*. Comparing root systems of healthy plants with diseased plants of the same age.

Sweet peas infested with *Thielavia* have a dwarfed and sickly appearance. The fungus does not seem to kill it, but merely to produce an arrested development. The infected plants are useless for commercial purposes, as they fail to set flowers.

Pathogenicity. Chittenden₂₀ seems to have been unable to infect healthy sweet pea seedlings with the fungus *Thielavia basicola* under normal conditions of growth. It was only when his plants were overwatered that the fungus became an active parasite.

In my own inoculation experiments, healthy sweet pea seedlings have been readily infected by placing a pure culture of the fungus on the roots of the plants growing in sterile soil. In two to three weeks the roots were thoroughly diseased. Overwatering was not found necessary to bring about infection, although such treatment as well as injury to the roots favor the fungus in its activity. Another method adapted for proving the pathogenicity of the fungus was to sow pure cultures of the fungus together with sterilized seed (seeds treated in a solution of formaldehyde, 5 parts in 100 of water for 1-2 hr.) in sterile pots and soil. Checks were also sown with sterilized seeds in sterile pots and soil but without the fungus. Six days after sowing both lots of seeds germinated and both check and infected seedlings apparently grew equally as well. Beginning with the third week, infected seedlings ceased growing, whereas the checks made considerable progress. After six weeks the infected seedlings were seen to be decidedly dwarfed and pale green in color reproducing the typical symptoms of the disease as observed in the field. The check seedlings have by this time made decided growth. An examination of the roots of the infected seedlings revealed a diseased condition as found in the field, namely, absence of a well developed root system, and a blackening of the affected parts. The infection experiments were repeated five times always with the same result. In no case was the *Thielavia* seen to kill the host, but in each case a dwarfed condition of the plant was the result.

Infection of Sweet Peas with Thielavia from Other Hosts. It was found desirable to determine whether there existed any racial strains or physiological species of *Thielavia basicola*. Accordingly, pure cultures of the fungus obtained from cowpea, violets, parsnip and tobacco, were inoculated on sweet peas, using the same method of inoculation as previously described. In connection with this experiment

a parallel series of inoculations was also run, using the *Thielavia basicola* from the sweet pea. The results obtained were the same, i. e. the *Thielavia* fungus when taken from other hosts than the sweet pea will readily infect the sweet pea, thus showing the absence of physiological strains or species.

Morphology of Thielavia basicola. Our studies and observations on this fungus have brought out facts found by other investigators. The mycelium of the fungus is hyalin, septate and branched. The hyphæ average from $3-4\mu$ in width. The mycelium becomes more or less greyish with age. There are three kinds of spore forms produced. 1. Endospores, so called because they are formed inside a special thread of the mycelium. This is the spore form that is most commonly met with in pure cultures of artificial media. The endo-




Fig. 3. Endospores.
 Figs. 4, 5. Chlamydospores breaking up into individual spores.
 Fig. 6. Chlamydospores, unbroken.
 Fig. 7. Ascospores.
 Fig. 8. Ascus.

spore case is formed on terminal branches. It has a somewhat swollen base and a long tapering cell (Fig. 3). The endospores are formed in the apex of this terminal cell and are pushed out of the ruptured end by the growth of the unfragmented protoplasm of the base. They are hyalin, thin-walled, oblong to linear $10-25\mu \times 4-5\mu$.

The second kind of spores formed are the chlamydospores (Figs. 4-6). These are thick-walled dark brown bodies, born on the same mycelium as the endospores, and average from $20-50\mu \times 10-18\mu$, and correspond to the *Torula* stage of Berkeley's classification.

The third kind of spores are the ascospores (Fig. 7.) These are lenticular in shape 12×5^u and are born in asci which in turn are born in spherical black perithecia. We have not as yet found the ascospore stage (Fig. 8) on the affected host, altho it is said to appear quite commonly on other hosts affected by this fungus.

Peglion₂₁ in 1897 was apparently the first to grow the fungus in pure culture. Aderhold₂₂ in 1903 reports to have grown the fungus in pure culture. Clinton₂₃ and Gilbert₂₄ experienced considerable difficulty in obtaining a pure culture of the fungus, due to the fact that the chlamydospores when taken from old diseased roots fail to germinate by being overrun with bacteria. At no time did the writer experience any difficulty in obtaining a pure culture of Thielavia from diseased sweet pea tissue. The following method was adapted from Manns₂₅. Portions of the diseased roots are placed in a test tube in a solution of 1-1000 HgCl₂ in a 50 per cent. alcohol and thoroughly shaken for thirty seconds. This will kill all surface contaminations. The disinfectant is then poured out and the material is rinsed three times in sterile water, the object being to remove all traces of mercuric chloride. Each tissue fragment is then taken separately and crushed with a sterile forceps in a tube of medium which has been melted and cooled to the proper temperature. The crushed tissue is now mixed with the medium in the tube, and the whole poured into a petri dish. After five or six days a pure growth of the fungus appears in the petri dish. The growth in this case resulted from the mycelium which has been crushed and liberated from the deeper tissue.

The fungus will grow on a variety of media. It grows well on sterilized vegetable plugs as those of potato, beet, carrot, sweet potato, corn stalks, and parsnip and particularly well on corn meal. Both endospores and chamydospores are produced on these media, but in no case did I obtain the perfect stage, although it was often looked for. Previous investigators too have never succeeded in obtaining the ascospore stage on pure culture.

Pathological Conditions of the Diseased Host. Reference has been previously made to the fact that the disease produced by Thielavia is confined to the root system. Infected plants have little or no root system at all, or if present it is charred and invaded by the fungus. The question arises, how do the plants persist such a long time without collapsing? It was observed that sweet peas affected with

Thielavia, constantly make an attempt to produce new roots, but as fast as they are formed they are invaded by the fungus. It is possible, therefore, that these new rootlets help the host to persist so long, and yet not long enough to enable it to make any growth. It is also probable that there is a close symbiosis between host and parasite with the latter getting the upper hand. As far as observed from cross sections of that part of the stem which lies closest to the roots, the fungus is seen to invade all parts of the tissue with the exception of the xylem vessels. This fact, that the fungus does not enter the conducting vessels permits an upward movement of the water, and this is sufficient to prevent the host from dying.

ROOT ROT, *Corticium vagum* B. & C.

Historical data, European literature. The first mention of Rhizoctonia can be traced to Duhamel₂₆ who in 1728 described a disease of Saffron (*Crocus sativus*). He considered the sclerotia to be a special plant and the hyphæ its roots, and named it Tuberoïdes. In 1782 Fougereux de Bondaroy₂₇ found asparagus plants which grew near a diseased saffron field to be likewise affected with Tuberoïdes. The first attempt to place the fungus in a systematic position was made by P. Builliard₂₈ who referred it to the Truffles-as Tuber parasiticum. Persoon₂₉ placed it in the Genus Sclerotium, and called it Sclerotium crocorum. De Candolle₃₀ was first to use the name Rhizoctonia. He at that time distinguished three species, *R. crocorum*, *R. medicaginis*, and *R. mali*. Nees₃₁ in 1817 refers to a fungus disease of the crocus which he calls Thanatophytum crocorum. From an examination of his figures there can be no doubt but that it is Rhizoctonia. In 1830, DUBY₃₂ described a fungus disease of *Allium ascalonicum* and named it Rhizoctonia allii. In 1843 Leveillé₃₃ describes Rhizoctonia as attacking *Rubia tinctorum*, *Solanum tuberosum*, *Phaseolus*, and other plants. (The species of Rhizoctonia is not stated.) In 1851 the Tulsane₃₄ Brothers placed all the known forms of Rhizoctonia in one species which they called Rhizoctonia violacea. However, from Kuhn's critical work a few years later, it seems advisable to maintain the distinction between *R. solani* and *R. medicaginis*. In Kukn's₃₅ work which was published in 1858, a brief account is given of the smooth sclerotia of *R. solani* contrasted with the wooly sclerotia of *R. medicaginis*. Distinction is also made of *R. crocorum* and *R. medicaginis*, the latter is stated to attack beets and carrots. In 1903

Erikson₃₆ concerns himself chiefly toward the discovery of biologic forms of the fungus.

In 1905, Gussow₃₇ described the disease on potato and lucern, and he considers *R. solani* and *R. violacea* as one and the same.

In 1912 Shaw₃₈ investigated the morphology and parasitism of *Rhizoctonia* with a view of obtaining a better understanding of the supposed different species. Shaw concludes that the name *R. violacea* should be retained for all the non-fruiting forms with macrosclerotia, and the name *Corticium vagum* be given to the fruiting stage of the macrosclerotia of *R. violacea*, while the forms with microsclerotia should be identified as *R. solani* Kuhn.

American Literature on Rhizoctonia. The American references to *Rhizoctonia* are as follows:

In 1891 Pammel₃₉ describes a rot of the beet root which he attributes to *Rhizoctonia betae* Kuhn.

In 1892 Atkinson₄₀ found a sterile fungus causing a damping-off of cotton, which he called "sore shin." The fungus can no doubt be referred to as *Rhizoctonia*. Later, Stone and Smith₄₁ give an account of a lettuce disease due to *Rhizoctonia*.

In 1901 Duggar and Stewart₄₂ describe extensively a list of hosts attacked by *Rhizoctonia*. However, the species of the fungus is not given.

In 1904 Rolfs₄₃ reports on a potato disease due to *Rhizoctonia*, in which he found the fertile stage *Corticium vagum* B. and C. Specimens were sent by Rolfs to Dr. E. A. Burt, who pronounced it a variety of *Corticium vagum* B. and C. and for which he has suggested the name *Corticium vagum* B. and C. var. *solani* Burt.

In 1909 Stevens and Hall₄₄ described a disease of the apple, pear and quince, which they attributed to a sterile fungus *Hypochnus ochroleuca* Noack. The fungus is described as having small sclerotia and there is no doubt but that it is a *Rhizoctonia*. Several workers have claimed to have connected different fruiting stages with that of *Rhizoctonia*. In 1869 Fuckel₄₅ stated that the Ascomycete *Byssothecium circinans* Fekl. was the perfect form of *Rhizoctonia*, both stages were found on decaying stems of *Medicago sativa*. Prunet₄₆ also observed this association of *Rhizoctonia* on lucern with an ascomycete.

Hartig₄₇ found a *Rosellinia* associated with a *Rhizoctonia* on the roots of oak.

Frank₄₈ reports *Rhizoctonia violacea* on grapes to be associated with *Thelephora*, which he named *Th. rhizoctoniae*. However, none of the above authors have carried on any cultural work to prove the validity of their claims, hence none of the statements can be accepted as valid.

Rolfs₄₉ found a basidiomycete associated with *Rhizoctonia* on potato. Pure cultures obtained from spores of the basidiomycete always gave a *Rhizoctonia*, thus proving definitely the relationship of the two forms. Rolfs basidiomycete is already known as *Corticium vagum* B. and C.

American, European and East Indian Hosts Subject to the Attacks of Rhizoctonia.

<i>Name of Host</i>	
Sugar beet	Duggar
bean, pea	" , Leveille
carrot	"
celery	"
cotton	"
lettuce	"
potato	" , Leveille, Rolfs.
crocus	Duhamel, Bulliard, Pers.
asparagus	Fougeroux
alium sp.	Duby
lucern	Kuhn
ground nut	Shaw
cowpea	"
jute	"
soy bean	"
mulberry	"
sweet pea	Taubenhaus

Rhizoctonia Attacking Sweet Peas. As far as could be ascertained no mention could be found in literature of a *Rhizoctonia* disease of sweet peas. I have observed it during winters of 1911 and 1912 on greenhouse specimens sent in by different sweet pea growers. In the fall of 1913, diseased sweet pea seedlings attacked with *Rhizoctonia* were collected in the greenhouse of the University of Pennsylvania. From correspondence with Plant Pathologists, A. D. Selby reports it in Ohio, W. G. Sackett in Colorado and E. C. Stackman in Minnesota. There seems no doubt but that the *Rhizoctonia* root rot of sweet pea is much more widespread than is reported.

Symptoms of the Disease. Severely infected plants have practically no root system (Fig. 9). In less infected plants only one or two rootlets may be destroyed. The fungus produces a browning effect of




Fig. 9. Root rot caused by *Rhizoctonia*. (A) healthy. (B) diseased.

the root before total destruction sets in. In very early stages of infection the seedlings are seen to have a wilted appearance; as the disease progresses the infected seedlings fall over and collapse. The fungus is not often confined to the roots alone. It is often seen to work its way up the stem and this may produce a constricted area which marks it off from the healthy part. The fungus being a soil organism, it is usually introduced with manure; infection can take place at any part

of the roots, or at the stem near the roots. When the latter is the case reddish sunken spots are observed at the base of the stem. It seems that only young seedlings can be quickly destroyed by the fungus whereas older plants seem to linger for considerable time altho such plants remain dwarfed, sickly looking and valueless for commercial purposes.

Pathogenicity. The pathogenicity of the sweet pea *Rhizoctonia* is readily proven by planting sterilized seeds in sterile soil and pots which were inoculated with a pure culture of the fungus. The best material to use is somewhat old cultures which have well developed sclerotia. It is from the latter that the fungus begins to vegetate and to spread in the soil. Five pots were inoculated with the fungus and two were kept as checks. The checks germinated and grew well, whereas none of the seeds germinated in the infected pots (Fig. 9a). In digging out some of these seeds they were found to be invaded with the fungus hyphæ of the *Rhizoctonia*. A pure culture may be readily obtained from these seeds, thus proving that the *Rhizoctonia* is a pathogenic organisms. Young seedlings may likewise be infected by the fungus, but as already indicated older plants are more resistant as they can live for some time with the fungus on them.

Morphology and Identity of the Sweet Pea Rhizoctonia. So far as my studies have gone, only two stages have been found of the sweet pea *Rhizoctonia*.

1. The *Rhizoctonia* stage.—This consists of long and narrow hyphal branches varying in color from hyalin to reddish brown (Fig. 10). These hyphæ are either aerial or are embedded in the substratum, varying according to the media on which it is grown. It is this hyphal growth which is most active in the parasitism of the fungus.

2. The Sclerotial stage.—In cultures which are from three to four weeks old numerous sclerotia are formed. These sclerotia are made up of closely interwoven short barrel-shaped hyphae (Fig. 11).

According to Shaw₅₀ *Rhizoctonia solani* Kuhn produces only microsclerotia while *Corticium vagum* B. and C. produces macrosclerotia. After repeated attempts the corticium stage of the sweet pea *Rhizoctonia* could not be obtained on artificial media. This accords with the findings of Shaw and Rolfs who could not obtain the perfect stage on culture media but found it several times on the affected host. However, as the sweet pea *Rhizoctonia* produces macrosclerotia and as




Fig. 9A. Root rot caused by *Rhizoctonia*. To the right the soil was inoculated with the fungus, resulting in no germination. At the left the soil was free from the fungus, resulting in good germination.

already pointed out by Shaw the macrosclerotia produce the *Corticium* stage: the sweet pea organism is therefore referred to as *Corticium vagum* B. and C.

Pathological Conditions of the Host. The *Rhizoctonia* fungus when attacking other hosts, is known to be confined primarily to the cambium layer of the plant. With the sweet pea, conditions are sim-

ilar. The fungus attacks the phloem of the bundles and makes its way into the parenchyma cells as well as to the epidermal cells. The effect produced is loss of turgidity, wilting, and early collapse of the




Fig. 10. Young hyphae of *Rhizoctonia* from sweet pea.

Fig. 11. Barrel shaped hyphae from sclerotium of the same fungus.

host. Infection may take place in the base of stem first and in this case the fungus invades both stem and root, or it may start at the roots first, then gradually work up to the stem. In either case death of the seedling is a natural consequence. In cases where the roots are first attacked by the fungus, the former deteriorates so rapidly that when pulling out a plant, it is found to be without any root system.

CHAETOMIUM ROOT ROT, *Chaetomium spirochaete* Patt.

Historical. In the fall of 1912 Prof. A. C. Beal of Cornell University sent me specimens of diseased sweet peas grown in the greenhouse for diagnosing the cause of the trouble. The disease was readily located in the roots. A fungus was found invading the interior tissue of the latter but no fruiting stage of any kind could be found which would help to identify the fungus. Crush cultures were made at once from the diseased tissue, the method employed being the same

as that described for *Thielavia basicola*. Some forty poured plates of nutrient agar were made in all. In 5 days a pure culture of a fungus appeared in all the plates with the exception of one which showed a *Fusarium*. The cultures were watched closely and in two weeks perithecia-like bodies developed in abundance but no spores were formed. The fungus proved to be an ascomycete belonging to the genus *Chaetomium*, and determined by Mrs. Flora Patterson as *C. spirochaete* Patt. In mid-winter of that same year, more diseased specimens were sent to me from a florist in Illinois. These specimens showed the same symptoms as those observed on Prof. Beal's material and in this case, too, the trouble was confined to the roots. As previously stated, no fruiting stage was found on the affected tissue. Cultures made from this material gave a pure growth of *Chaetomium spirochaete*. A search through the literature failed to show the record of any of the known *Chaetomiums* to be parasitic on living plants. It is known for instance that *C. arachnoides* Masee, *C. simile* Masee, *C. bostrychoides* Zopf and *C. murorum* Cda., all grow on dung of various animals.

Reinke and Berthold₅₁ in their studies of the fungous diseases of the potato report to have found *Chaetomium bostrychoides* Zopf and *C. crispatum* Fekl. growing on rotted tubers. The above authors state that when germinated spores are placed on a cut surface of a healthy tuber they fail to penetrate the same, indicating the saprophytic nature of the fungus. Its presence on the decayed potatoes must have been secondary. The present thesis gives the first record of the parasitic nature of *Chaetomium*.


Pathogenicity. The fact that a pure culture of *C. spirochaete* was obtained from numerous platings of diseased material obtained from two different states at once led to the supposition that the organism was the cause of the disease. In order, therefore, to test out the pathogenicity of the fungus, the following experiments were tried. Out of ten sterilized pots and soil, five were sown with sterilized seeds (these were soaked in a solution of formaldehyde, 5 parts in 100 of water for one-half hour) inoculated with a pure culture of the fungus broken up in sterilized water. In the remaining five pots sterilized seeds were sown without the fungus to serve as checks. Both lots of seeds germinated and the seedlings of both the inoculated and the check pots seemed to grow well for about three weeks. After that time

the seedlings in the inoculated lots appeared to lose their green color and to become paler and yellow. The infected plants could be readily pulled out from the soil and the rootlets appeared to be half rotted by the fungus, whereas the check seedlings did not exhibit such symptoms. Cultures made by surface sterilizing the affected rootlets readily gave pure cultures of *Chaetomium spirochaete*. The experiment was repeated once more and this time, both checks and inoculated seedlings were watered frequently, care being taken to keep the soil very moist or even wet. This was accomplished by allowing the dish to remain filled with water in which the pots stood. In this case the checks again remained healthy, but after three weeks the inoculated seedlings had their roots mostly destroyed by the fungus; the infected seedlings could be readily pulled out from the soil. This time the greatest part of the root system was destroyed. Cultures made from parts of the remaining infected roots readily gave pure cultures of the fungus.

From these experiments it is shown that *Chaetomium spirochaete* altho perhaps a saprophyte will, under certain conditions, assume a parasitic nature on sweet peas. It was further shown that in poorly drained soils the virulent nature of the organisms becomes more pronounced.

Morphology and Physiology of the Fungus. The mycelium of the fungus is hyalin, closely septate and branched (Fig. 12) when grown in the substratum of the media. The aerial mycelium consists of long unbranched filaments and vary in color from very light to deep lemon. This seems to be produced within the fungous hyphae and later the yellow color is also transmitted to the media in which it grows.

Reinke and Berthold₅₂ report to have found a conidial stage connected with *C. crispatum*. In our work we have as yet not found any conidial stage of *C. Spirochaete*. As previously stated, we have not found any fruiting stages of the fungus on the affected host. In pure culture in artificial media perithecia appear in two weeks from the time of planting and in three weeks mature asci with spores are also formed. The perithecia are covered with darkish brown hair-like appendages, thus giving it a bristly appearance. The hairs are coiled at the apex and septate at different intervals; they are covered with very minute pointed warts (Fig. 13). The asci are very evanescent



Figs. 12-15. Showing (12) mycelium of *Chaetomium spirochaete*. 13, hairs. 14 and 15, asci. 16, ascospores.

and can only be seen in young cultures. In old cultures the ascus wall readily breaks so that it is difficult to make out the arrangement of the ascospores. There are 8 ascospores to an ascus (Fig. 14-15). The ascospores are apiculate (Fig. 16) at both ends. The wall of the ascospore is smooth, light brown when young and dark when old. The ascospores readily germinate in a sweet pea broth which is made up as follows:

Take 15 grams of ground sweet pea seeds to 1000 cc of water. Bring to a boil, filter, then add 15 grams of agar; bring to a boil, then filter, tube and sterilize in the autoclave for 15 minutes at 15 pounds pressure.

FUSARIUM ROOT ROT, *Fusarium lathyri* n. sp.

Historical. It seems that Tulasne₅₃ was the first to recognize the parasitic nature of *Fusarium*. In 1883 Hansen₅₄ describes a disease on oats which is attributed to *Fusarium gramineum* Corda. In 1884 Worthington Smith₅₅ describes a wheat disease due to *Fusispor-*

ium (*Fusarium*) *culmorum* W. Sm., and another disease on barley due to *Fusarium hordei* W. Sm. In 1889 E. F. Smith₅₆ mentions different related *Fusaria* isolated from the soil which play the role of plant pathogens. The species of the *Fusarium* are not given. In the same year, in their excellent report on the loose smut of wheat, Kellerman & Swingle₅₇ report on a *Fusarium* which lives as a parasite on the loose smut of wheat and which they named *Fusarium ustilaginis* K. & Sw. In 1892 Frank₅₈ reports *Fusarium heterosporium* Nees, a parasite on several graminaceous hosts. In the same year Atkinson₅₉ reports on a cotton disease due to *Fusarium vasinfectum*. In 1893 Rostrup₆₀ described an oat disease due to *Fusarium avenaceum*. In 1899, Smith₆₁ describes a disease of cotton, watermelon, cowpeas, and melons as due to a *Fusarium* whose perfect stage was believed to be *Neocosmopora vasinfecta*. In 1899 Woods₆₂ reports on a disease of Chinese asters due to a *Fusarium*. In 1900 Manguin₆₃ reports on the parasitic nature of *Fusarium roseum*. In 1901 Prillieux and Delacroix₆₄ reported on a carnation disease due to *F. dianthi*. In 1901 Bolley₆₅ reports on a flax disease due to *Fusarium lini*. In 1901 Sorauer₆₆ reports on a rye disease due to *Fusarium nivale*. In the same year Pammel₆₇ reports on a wheat disease due to *Fusarium roseum* Lk. In 1902 Smith₆₈ reports extensively on a wilt of Chinese Aster the same disease as previously reported by Woods₆₉. In 1902 Hennings₇₀ describes a disease on the black locust which he attributes to *Fusarium vogelii*. In 1903 Van Hall₇₁ describes a pea disease which he attributes to *Fusarium vasinfectum* Atk. var. *Pisi*. In 1904 Smith and Swingle₇₂ reports on the dry rot of potatoes due to *Fusarium oxysporum*. In the same year Osterwalder₇₃ reports on a fruit rot due to *F. putrefaciens*. In 1905 Owen₇₄ reports on a tomato disease which he attributes to *Fusarium erubescens* Appel. and v. Oven. This fungus is claimed to be different from *Fusarium solani*, *F. putrefaciens* and *F. Lycopersici*. In 1906 Appel and Schikarra₇₅ report on different species of *Fusaria* which induce disease in plants. In the same year Heald₇₆ reports on a bud rot of carnations due to a species of *Fusarium*. In 1906 Hedgcock₇₇ in his extended studies of chromogenic fungi reports *Fusarium roseum* as capable of discoloring wood. In 1907 Chiffot₇₈ reports on a pelargonium disease due to *Fusarium pelargonii*. In 1909 in their extended studies on corn rots, Burrill and Barrett₇₉ report on three species of *Fusaria* which attack the ear of corn. In 1910 Wolf₈₀ reports on a pansy disease as due to *Fusarium*. In the same year

Smith₈₁ reports on a banana disease due to *Fusarium*, and Cook₈₂ reports on the double blossom due to *Fusarium rubi*. In 1911 Bubak₈₃ describes a rot on ears of corn due to *Fusarium maydiperdum*. In 1912 Gifford₈₄ reports extensively on a damping off disease of coniferous seedlings due to *Fusarium*, the species not stated. There are of course many more *Fusaria* described, but the aim in this brief historical sketch is to mention the literature which bears more or less directly on the parasitic nature of *Fusarium*. In a very recent paper Wollenweber₈₅ describes several species of *Fusaria* parasitic on the potato. He also emphasizes the importance of morphological studies as well as infection experiments as the basis of classification in *Fusaria*.

Fusarium Root Rot of Sweet Pea. There is no record in the literature of a *Fusarium* disease of the sweet pea. Several complaints from florists have shown that they could not grow sweet peas under greenhouse conditions because of a root rot which developed early and in some cases destroyed the entire planting. Cultures made from the infected material or from the infected soil, and from seedlings sown in the laboratory on the infected soil, gave in each case a pure culture of *Fusarium*.

Symptoms. The first symptom of the disease is a sudden flagging of the leaves accompanied by general wilting and collapse of the seedling. Usually upon sowing the seeds a fair percentage germinate and reach the height of about 8 to 10 inches when they are attacked by the fungus. If the collapsed seedlings are allowed to remain on the ground, the stems will soon be covered with the sickle shaped spores. Eventually the decayed tissue rots and disintegrates and is soon invaded by small fruit flies which now begin to distribute the fungus from place to place by carrying its spores.

Pathogenicity. The pathogenicity of this fungus is readily proven by inoculating with a pure culture of the organism sterilized seeds planted in sterile soil. The seed germinate and reach a height of 7 to 8 inches but soon succumb to the attacks of the fungus. The fungus can be reisolated from the artificially infected seedlings and the disease induced at will (Fig. 17). The checks remain healthy provided of course all means of contamination are guarded against.


Morphology of the Fungus. The mycelium of the fungus is hyalin, septate and branched. At an early age the hyphae begin to



Fig. 17. *Fusarium* wilt or root rot. At left, healthy; at right, infected.

form chlamydospores. These are round hyalin bodies often filled with oil globules and are formed in the center of the hypha (Fig. 18), in this case the contents of the former collect into the chlamydospores. Usually also the chlamydospores are born at the tip end of the hyphae in chains of twos, threes and even fours (Figs. 19-22). Old cultures are practically one mass of chlamydospores. There are also two spore forms present and these appear as early as the third day in the pure culture. These are microconidia which are fairly abundant and macroconidia, varying from two celled to four celled. The average form is the three celled. Both micro- and macro- conidia are hyalin and smooth (Figs. 23-31). In old cultures the macroconidia shrink so that the septa become slightly pronounced (Figs. 25, 28-29). These old macroconidia soon lose their protoplasm or the latter breaks up

presenting a granular appearance. In young cultures the outer wall of the chlamydospore is smooth, but in old cultures it becomes slightly warty or covered with minute points (Fig. 19). No perfect stage has been found to accompany this fungus either in pure culture or on the host.



Figs. 18-31. *Fusarium lathyri*, showing chlamydospores and conidia.

Identity of the Fungus. There is no doubt but that the fungus belongs to the genus *Fusarium*. It produces its micro- and macrospores (sickle shaped) as well as chlamydospores which according to Wollenweber⁸⁶ are true characteristics of the genus *Fusarium*. The fungus has been grown in pure culture (Fig. 32) and on different media for two years and no perfect stage has ever appeared. Unless further studies prove differently it seems that the present *Fusarium* is a new species and the name *Fusarium lathyri* n. sp. is tentatively given to it. A description of the fungus follows:

Sporodochia slightly erumpent to superficial on the host, but not always present on culture media. Macroconidia sickle-shaped, slightly curved and fitting into the section Martiella of Wollenweber, 2 to 4 septate, the majority being three septate, 15.8×4.2^u — 30.8×5.6^u . Microconidia elliptical to oval 9.8×2.8^u — 14×3.50^u . Chlamydospores spherical, thick walled and spinulate when old, 7.3^u — 9^u borne singly or in chains of twos, threes and sometimes in fours.

ROOT ROT OR EEL WORM, *Heterodera radicolica* (Greef) Muller

Altho not belonging to the domain of this thesis the eel worm is here considered, first because of the important root knot disease it

produces, and the second, because it opens the way to several fungous parasites.

Heterodera radicola attacking many hosts. The eel worm is of an omnivorous nature. Marcinowski₈₇ gives a list of 235 species of plants affected with the pest. Almost all of the more important families of flowering plants are present in the list, as well as one gymnosperm and a fern. The plants include both Monocotyledons and dicotyledons, herbs, woody plants, annuals and perennials. Many of our garden plants and field crops are attacked by the pest.

Of the plants said not to be affected by *Heterodera*, Bessey₈₈ cites the following hosts: Crab grass (*Syntherisma sanguinalis*), redtop (*Agrostis alba*), Johnson grass (*Andropogon halepensis*), some varieties of oats (*Avena sativa*), *Bromus schraderi*, *Eustachys petrea*,




Fig. 32. Pure culture of *Fusarium lathyri*. the cause of sweet pea wilt

some varieties of barley (*Hordeum vulgare*), perennial rye grass, *Lolium perenne*, *Echinochloa frumentacea*, *Panicum miliaceum*, *Pennisetum* sp. timothy (*Phleum pratense*), rye (*Secale cereale*), *Andropogon sorghum*, *Triticum*, maize (*Zea mays*), *Euchlaena luxurians*, *Bidens leucantha*, *B. bipinnata*, *Gnaphalium purpurem*, *Helenium tenuifolium*, some species of *Solidago* and *Zinnia*.

Heterodera attacking sweet peas. In his excellent work on the root knot Bessey₈₉ mentions the sweet pea as attacked by *Heterodera radiculicola*. Chittenden₉₀ found the eel worm associated with *Thielavia* root rot.

In my own investigations I have found the eel worm capable of producing the root knot. I have often found this pest to be associated with *Rhizoctonia*. It seems very probable that the *Heterodera* worm in the case of the sweet pea opens the way to the attacks of *Rhizoctonia* and several other fungi. *Heterodera radiculicola* often produces the greatest amount of damage in light sandy soil, but seems unable to thrive in heavy clay soils. In every case where seen by the writer or where reported by florists and seedsmen the eel worm was most troublesome where sweet peas have been grown on too light soils in the greenhouse. No sweet peas have been reported to be attacked by the eel worm out of doors.

Symptoms. The disease is characterized by swellings on the roots. These are either small swellings formed singly, in pairs, or in strings, thus giving the affected root a beaded condition (Fig. 33) or the swellings may be very large so as to be mistaken for root nodules. However, these galls cannot be mistaken for the normal root nodules, because the latter are lobed and are attached at one end (Fig. 34), whereas the root gall produces a swelling of the entire surface of the part affected. Infected plants usually linger for a long time, but they can be distinguished by a thin growth and yellow sickly looking leaves and stems.

B. Diseases of the Aerial Parts of the Plant

STEM OR COLLAR ROT, *Sclerotinia libertiana* Fckl.

Synonymy.

Sclerotinia libertiana Fckl.

Peziza sclerotiorum Lib.

Sclerotinia libertiana Fckl.

“ *Kaufmanniana*


“ *Postuma* B. & W.

“ *Coemansii* Kick.

“ *sclerotiorum* Br.

Sclerotinia sclerotiorum Ad.

History of Sclerotinia libertiana Fckl. as a Plant Disease Producer. In 1860 Coemans₉₁ was the first to record a disease of carrots and turnip which he attributed to *Sclerotinia libertiana* (*Peziza sclerotium*). In 1886 De Bary₉₂ also reports on a disease of turnips



33



34

Fig. 33. Sweet pea roots affected with eel worm.

Fig. 34. Sweet pea nodules formed by nitrogen fixing bacteria.

and carrots due to *S. libertiana*. In 1887 Cohn₉₃ reports on a potato disease due to the same fungus. Smith₉₄ in 1890 reports on a hollyhock disease due to *Sclerotiana libertiana*. In 1891 Behrens₉₅ reports on a hemp disease due to this fungus. In 1892 Humphrey₉₆ reports

on a lettuce disease which he thought was due to *Botrytis cinerea*, the then supposed conidial stage of *Sclerotinia libertiana*. Humphrey again in 1893 reports on a cucumber disease due to *Sclerotinia libertiana*. In 1897 Prillieux and Delacroix₉₇ report on an important mulberry disease due to this same fungus. In 1900 in his extended studies on *Sclerotinia*, Smith₉₈ reports on the omnivorous nature of *S. libertiana*. Hedgcock₉₉ in 1905 reports on a serious disease of cabbage and cauliflower due to *S. libertiana*. In the same year Parisot₁₀₀ reports on a disease of Jerusalem artichokes in the western part of France and due to this same fungus. He also mentions the potato, bean, corn, carrot, turnip, rutabaga and flax as all being subject to the same disease. In 1906 Appel and Bruch₁₀₁ also report on a similar disease of turnip and parsnips. Masseron₁₀₂ in 1907 reports this fungus to produce a serious disease on the garden and field pea. In 1911 Westerdijk₁₀₃ reports a wide range of hosts to be attacked by the fungus such as rape, cabbage, cauliflower, kohlrabi, sesame, bean, pea, vetch, clover, lettuce, Jerusalem artichoke, dahlia, zinnia, sugar beets, carrots, turnips, parsnips and beets.

Sclerotinia libertiana Fckl., a Fungous Disease of the Sweet Pea. As far as I could ascertain there is no mention in literature of *S. libertiana* attacking the sweet pea. I have first noticed this disease on greenhouse specimens sent in by a grower in Pennsylvania. My first record of the disease appeared in the Florist Exchange₁₀₄. Observations so far seem to show that the disease is limited to sweet peas under greenhouse conditions only. A special effort was made to find this disease out of doors but without success. It is well known that under certain conditions unfavorable to the host the fungus can attack a variety of hosts which grow in the open. That the fungus has not been found so far to attack sweet peas out of doors does not limit its appearance in the field at any time that climatic conditions are unfavorable to the host.

Symptoms. This is usually a seedling disease (Figs. 35-36) altho it may attack plants of all ages (Fig. 36).* And it is most severe in poorly ventilated houses or in beds which are overwatered and which lack proper drainage. The disease when present does its work quickly. Affected plants first show a wilting of the tip and flagging

*Figs. 36, 41. Electrottype, Florists' Exchange. Photographs by the author.

of the leaves and then the seedlings fall over and collapse. The fungus *Sclerotinia libertiana*, altho a soil organism cannot attack the roots of its host. The fungus penetrates the collar of the stem and completely invades the vessels of the host, thus checking the upward flow of the water from the roots. Freshly collapsed plants have a watersoaked



Fig. 35. *Sclerotinia* wilt of sweet pea seedlings. (a) healthy plants. (b and c) infected seedlings.

appearance, later to be overrun with a white web of fungus mycelium, finally to be followed by sclerotia which are found here and there on or within the affected stems.

Pathogenicity. In order to establish definitely the relationship of the fungus to this disease of sweet peas under glass, sterilized seeds were planted in sterilized pots and soil in the laboratory. All the seeds germinated and the plants were allowed to grow for three weeks, in a perfectly healthy state. Then the pots were divided in



Fig. 56. Damping off caused by sclerotinia.

two lots; one lot was left as a check and the other lot was inoculated with the pure culture of the *Sclerotinia* by introducing pieces of the fungus in the soil. Both lots, check and infected plants, were covered with bell jars to imitate the moisture conditions of the greenhouse.

After four to six days, wilting of the inoculated seedlings began, whereas the checks remained healthy. This was repeated several times with always the same results. This conclusively shows that the fungus *Sclerotinia libertiana* is able, when present in the soil, to produce a disease on sweet peas under glass. The fungus is usually brought into the greenhouse with the soil, or with the manure. Cross inoculation with the fungus from the sweet pea, lettuce, turnip and cucumber on the sweet pea produced typical cases of wilt, thus proving conclusively that the fungus from the sweet pea is the same as the *Sclerotinia libertiana* of the hosts mentioned above.

Identity of the Fungus. The fungus from the sweet pea was run on artificial media with parallel cultures of *S. libertiana* from lettuce, cowpea and cucumber. There was no difference observed in the manner of growth nor of sclerotial formation of these strains. Under pathogenicity, I have shown that cultures of *S. libertiana* obtained from hosts such as lettuce, turnip and cucumber readily produced the typical wilt of the sweet pea similar to that obtained when inoculations are made with the fungus from the sweet pea on the sweet pea. In order to further verify the identity of the fungus, sclerotia from cultures three months old were placed in small flat covered chambers containing sterile moist sand. These were placed outdoors in the cold for four weeks, then brought in the laboratory and kept at room temperature. The moisture was maintained by adding every three to four days some sterile water. In nine weeks the sclerotia germinated by sending out from each a number of grayish stalks, and in two weeks the apothecia developed at the tip of the stalks. In shape and measurement of asci and ascospores the fungus answered in every description that of *Sclerotinia libertiana*.

Morphology of the Fungus. In my work I find no conidial stage of a *Botrytis* type or of any other type to be connected with *Sclerotinia libertiana* of the sweet pea. This is in accord with the studies made by Smith, R. E.¹⁰⁵ There seems no doubt but that *Botrytis cinerea* which is often found to accompany *S. libertiana* is in no way connected with the former.

POWDERY MILDEW, *Erysiphe polygoni*?

The sweet pea mildew was first described by Masseur¹⁰⁶ as being prevalent in England. *Erysiphe polygoni* was attributed as the cause, both of sweet pea mildew and that of the edible

garden pea. In this country Stewart¹⁰⁷ was the first to record a powdery mildew on the sweet pea. However, Stewart did not find the perithecial stage, and hence the fungus was not determined. The powdery mildew is a very prevalent disease on




Fig. 37. Anthracnose disease of sweet pea on stem and peduncles.

greenhouse sweet peas, and on irrigated fields or where they are planted on a large scale for seeds. Ordinarily, however, in small garden lots, and especially where the plants do not receive any water, the disease is practically unimportant since the attack is usually very mild during the active growing season, but becomes somewhat more abundant when the plants have passed all usefulness. The writer had the opportunity of collecting specimens at random from six large houses, and from three acres of out-door sweet peas in Mass. and from a similar three acre plantation in Pa. Like Prof. Stewart, the writer has only met with the conidial or *Oidium* stage. On our own




Fig. 38. Anthracnose disease of pods and seeds.

sweet pea field, we have carefully watched for a perithecial stage but without success. Late in the fall, badly infected leaves have been collected and put away to winter over, but that material up to date, April, 1913, has failed to develop perithecia.

ANTHRACNOSE, *Glomerella rufomaculans* (Berk.) S. & V. Sch.

A very serious anthracnose disease of the sweet pea on the Delaware Experiment Station farm was called to the attention of the writer during the latter part of July, 1910. This disease proved to be the same or very similar to the one reported by Sheldon₁₀₈ from West




Fig. 39. Anthracnose disease affecting the sweet pea leaf.

Virginia in 1905, and is very probably the so-called "wilt" which has been so often referred to in old seed catalogues and treatises on sweet peas.

Symptoms. The disease occurs on the stems, leaves (Fig. 39), flowers and pods but is most severe on the latter. There is a general wilting of the affected parts followed by dying, which begins at the tips of the younger shoots and works downward (Fig. 37). The older parts of the plants are not killed immediately but may persist for some time after being attacked by the fungus. The dead parts shrivel, become brittle, and are soon covered with minute acervuli. The affected pods are at first a dirty white in appearance but assume a dull color, which is due to the presence of the acervuli. A definite canker, which is so characteristic of the bean anthracnose is not produced. Although the disease on the stems seems to be restricted to the youngest growths, the pods may be infected at any stage of their development. The seeds of the diseased pods are always infected, become shrivelled and frequently do not reach maturity (Fig. 38.)

Pathogenicity. Sheldon called attention to the identity of the *Gloeosporium* of the sweet pea with *Glomerella rufomaculans* (Berk.) Spauld. & von Sch. which is the cause of the bitter rot of the apple. With this in mind, the writer made the following experiments in the autumn of 1910. Sweet pea seeds, which to all appearances were perfectly healthy, were carefully selected, sterilized by immersion for 15 minutes in a 5% solution of formaldehyde, and planted in pots in soil which had been sterilized by heating for one hour in the autoclave. The seeds germinated in 5 days and the seedlings were allowed to grow for three weeks. Fifty plants were allowed to grow in each pot. The temperature of the room ranged as high as 72° F. during the day and several degrees lower during the night, but not low enough to injure the plants. All the seedlings grew well and were perfectly healthy. The day before inoculation the pots with the seedlings were covered with bell jars thus forming moist chambers. These covers were removed one day after inoculation. Two methods of inoculation were employed: (1) the introduction of spores into the living stems through punctures made with a sterilized needle; (2) by liberally spraying the surface of the plants by means of an atomizer with spores suspended in sterilized water. Fruits of apples and pears on the trees in the orchard were also treated in the following manner: Healthy fruits on the trees were first washed with a 5% solution of formaldehyde and then rinsed with distilled water. They were then inoculated through sterile needle punctures and covered with paper bags. For the inoculation, pure cultures were used of *Gloeosporium*

from various sources as indicated below. The results of these experiments are given in Table I.

The data in Table I show: (1) that the original organism of the sweet pea is pathogenic to the sweet pea, and also to the apple in which

TABLE I

Source of Gloeosporium culture	Number and kind of plants inoculated	Method of inoculation	Date of inoculation	Results of inoculation	Checks
Sweet Pea	50 sweet pea seedlings, 3 weeks old	Puncture	Sept. 26	Oct. 2, all dead	50 all healthy
" "	Same	Atomizer	Oct. 26	Nov. 7, " "	" "
" "	Same	"	Nov. 14	Nov. 21, " "	" "
Apple	Same	Puncture	Oct. 10	Oct. 18, " "	" "
"	50 sweet pea seedlings 4 weeks old	"	" "	Oct. 22, " "	" "
"	Same	Atomizer	Nov. 20	Nov. 29, 44 "	" "
"	Same	Puncture	Oct. 26	Nov. 1, 31 "	" "
"	Same	"	Nov. 14	Nov. 20, all "	" "
"	Same	Atomizer	Oct. 5	Oct. 14, 37 "	" "
"	50 sweet pea seedlings 8 weeks old	"	Oct. 7	Oct. 17, 42 "	" "
Sweet Pea	12 apples on tree	Puncture	Oct. 7	Oct. 17, typical bitter rot	5 "
" "	16 " "	"	Oct. 8	" " " "	12 "
Apple	15 " "	"	" "	" " " "	16 "
"	21 pears on tree	"	" "	" " " "	8 "
Sweet Pea	18 " "	"	" "	" " " "	11 "
" "	16 " "	"	" "	" " " "	12 "

it causes the typical bitter rot; (2) that *Gloeosporium fructigenum* Berk. from the apple causes a disease on the sweet pea which is similar to the disease caused by the original sweet pea *Gloeosporium*. This definitely proves that *Glomerella rufomaculans* (Berk.) Spauld. & von Sch. is the cause of the anthracnose disease of the sweet pea.

Relation of other Gloeosporiums to the Sweet Pea Disease.

While working on the question of the identity of the bitter rot of the apple and the anthracnose of the sweet pea, it was considered desirable to determine whether other species of *Gloeosporium* could produce an anthracnose of the sweet pea similar to that caused by the bitter rot organisms of the apple. Therefore, sweet pea seedlings were inoculated with spores from pure cultures of five different *Gloeosporium* then in stock in the laboratory. The results of these experiments are given in Table II.

TABLE II

Species of Gloeosporium	Number and kind of plants inoculated	Method of inoculation	Date of inoculation	Results of inoculation	Check
Gloeosporium gallarum	50 sweet pea seedlings	Puncture	Oct. 26	Nov. 7, 42 dead	50 all healthy
"	"	"	Nov. 14	Nov. 25, 41 dead	"
"	"	Atomizer	Nov. 30	Dec. 12, 8 dead	"
"	"	"	Dec. 2	Dec. 14, 19 dead	"
"	"	"	Oct. 26	Nov. 7, 34 dead	"
Gloeosporium officinale	"	Puncture	Nov. 14	Nov. 25, 43 dead	"
"	"	"	Nov. 30	Dec. 12, 50 dead	"
"	"	Atomizer	Dec. 2	Dec. 14, 50 dead	"
"	"	"	Oct. 26	Nov. 7, 50 dead	"
Species from May apple	"	Puncture	Nov. 14	Nov. 26, 42 dead	"
"	"	"	Nov. 30	Dec. 12, 17 dead	"
"	"	Atomizer	Dec. 2	Dec. 14, 19 dead	"
"	"	"	Dec. 26	"	"
Glomerella psidii	"	Puncture	"	Nov. 20 failure	"
"	"	Atomizer	"	"	"
Species from Persea	"	"	Oct. 20	"	"
"	"	Puncture	"	"	"
Gloeosporium gallarum	15 pears on tree	"	"	Nov. 15, typical bitter rot	7 "
Gloeosporium Officinale	12 "	"	"	"	5 "
Species from May apple	14 "	"	"	"	9 "
Glomerella psidii	19 "	"	"	Nov. 15, not typical bitter rot	6 "
Species from Persea	16 "	"	Oct. 5	"	6 "
Species from sweet pea	106 pole lima on pods	"	Oct. 7	Oct. 19, successful	16 "
Species from apple	66 "	"	"	"	12 "
Species from May apple	24 "	"	"	"	9 "
Gloeosporium Officinale	32 "	"	"	"	"
Gloeosporium gallarum	42 "	"	"	"	"
Glomerella psidii	60 bush lima bean	"	"	"	"
Species from sweet pea	42 "	"	"	"	19 "
Species from apple	49 "	"	"	"	12 "
Species from apple	40 "	Atomizer	"	Oct. 26, failure	12 "
Species from sweet pea	58 "	"	"	"	12 "

From Table II it will be seen that the Gloeosporium from the May apple fruit (*Podophyllum peltatum*), *G. gallarum* Ch. Rich. from oak gall, and *G. officinale* E. & E. from sassafras leaves, are able to infect sweet pea seedlings through puncture inoculations as readily as the sweet pea or the apple Gloeosporium. Furthermore, that *G. officinale* readily infects sweet pea seedlings by atomizer inoculations, the infection being nearly 100%. While *G. gallarum* and the species of Gloeosporium from the May apple fruit also infect sweet pea seedlings, the percentage of successfully inoculated seedlings is smaller with the atomizer inoculations than when the inoculations are made by needle punctures.

Glomerella psidii (G. Del.) Sheldon and the Gloeosporium from *Persea* failed to infect sweet pea seedlings.

The apple trees in the old orchard of the Experiment Station did not bear enough fruits to permit inoculation experiments with the above five organisms; hence Kieffer pear trees which bore heavily were chosen for this purpose. They were accordingly inoculated with the five Gloeosporiums already mentioned. The results of these inoculations (Table II), show that the species of Gloeosporium from the May apple *G. gallarum* and *G. officinale* produce the typical bitter rot on the pear, while the Gloeosporium from the guava and *Persea* infect the pear, but cause dull spots in which the acervuli are black and the spores are borne on long black conidiophores. Similar results were obtained when pear fruits were inoculated with the same Gloeosporium and kept in moist chambers in the laboratory. These experiments also show (Table II) that all the Gloeosporiums here considered, except the species from guava and *Persea* very readily produced an anthracnose disease on the pods of the lima beans in the field, which was similar to the anthracnose of the sweet pea, but quite unlike the bean anthracnose, *Colletotrichum lindemuthianum* (Sacc. & Mag.) B. & C. All the Gloeosporiums referred to above attacked the pods of the lima beans in the field when the inoculations were made by means of punctures, but not otherwise. The spots produced on the lima bean pods by *Glomerella psidii* are gray with grayish acervuli and made up of black setae very similar to those of a true *Colletotrichum*, but unlike *C. lindemuthianum*. None of the above species of Gloeosporiums would infect bean or vetch seedlings. The same precautions were taken in inoculating the bean pods and pears in the field as with the apples.

Further Studies of some Gloeosporiums and their relation to the Sweet Pea Anthracnose. The first year (1910) the inoculations were carried on at a time when both the apples and the pears were almost mature, and ripe fruits being a more favorable medium, since they are physiologically less active than young ones, it was felt advisable to start the present inoculations of the apples and pears in the orchard at a much earlier date, this time using more organisms. The inoculations were begun in 1911 when the fruits were the size of a grape, and were repeated at various stages of their development. The Kieffer pear and the Rubicon and Paradise Sweet apples were selected for this purpose. The inoculations were made by means of punctures through the cuticle. For each organism a different sterilized needle was used. Natural infection of the inoculated fruits was prevented by means of heavy paper bags which were tied to the limbs to which the inoculated fruits were attached. Any inoculated fruits which happened to drop off fell into the bags and were retained there. In every case where infection occurred it first appeared at the point of inoculation. For each organism eight fruits were used as checks. These were punctured with a sterile needle and covered with paper bags, and in all cases remained healthy. Investigations were also carried on with sweet pea, specimens of which were grown in the laboratory from carefully selected and sterilized seeds grown in sterilized pots and soil. Checks were also used, fifty seedlings for each organism, and these in all cases remained healthy, although they were punctured with a sterilized needle. Only spores from pure cultures were used for the inoculations. The results obtained are given in Table III.

From Table III it will be seen that *Glom. rufomaculans* from apple and sweet pea, *Gloe. gallarum* Ch. Rich., *Glom. gossypii* (South) Edg., *Gloe. diospyri* E. & E., *Colletotrichum phomoides* (Sacc.) Chest., and *C. nigrum* Ell. & Halst. produce the typical anthracnose disease on the sweet pea and the symptoms produced by all the above organisms were identical with those produced by the original Gloeosporium isolated from diseased sweet pea plants in the field. Many more inoculations than are indicated in Table IV were made with the above named organisms on the sweet pea. They were omitted from the table, since the results obtained were similar to those given above. The data in Table III further show that *Gloe. piperatum* E. & E. failed to infect the sweet pea by atomizer inoculation, while infection by puncture inoculation was fairly successful. When this organisms

was reisolated from seedlings infected by puncture, it regained its virulence, and then became able to infect sweet pea seedlings by atomizer inoculation. *Glom. rufomaculans* from the fig failed to infect the sweet pea, and, as will be seen later, it also failed to infect apples and

TABLE III

Species of fungus	Methods of inoculation	Date of inoculation	Results of inoculation
<i>Glomerella rufomaculans</i> from apple.....	Atomizer	June 16	June 28, 99% infection
<i>Glom. rufomaculans</i> from sweet pea.....	"	May 21	June 2, 92% "
"	"	May 25	June 6, 90% "
"	"	June 16	June 21, 95% "
"	"	June 29	July 11, 41% "
"	"	July 27	Aug. 10, 80% "
"	"	Nov. 1	Nov. 10, 100% "
<i>Gloeosporium gallarum</i>	"	June 16	June 28, 91% "
<i>Gloe. piperatum</i>	"	May 25	June 12, no "
"	Puncture	"	June 12, 2% "
"	"	June 29	July 6, 41% "
"	Atomizer	June 4	June 21, no "
" *	"	July 27	Aug. 9, 50% "
<i>Glom. gossypii</i>	"	May 21	June 2, 82% "
"	"	May 25	June 12, 60% "
"	"	June 21	July 7, 100% "
"	"	June 29	July 7, 80% "
<i>Gloe. diospyri</i>	"	Oct. 28	Nov. 10, 82% "
<i>Colletotrichum phomoides</i>	"	"	Nov. 10, 80% "
<i>C. nigrum</i>	"	"	Nov. 10, 92% "
<i>C. gloeosporioides</i>	"	July 3	July 14, no "
"	Puncture	Aug. 1	Aug. 10, " "
<i>Glom. rufomaculans</i> from fig.....	"	May 21	June 2, " "
"	Atomizer	"	June 2, " "
<i>Gloeosporium</i> from <i>Populus deltoides</i>	"	June 29	July 11, " "
"	Puncture	May 25	June 12, " "
<i>C. lindemuthianum</i>	"	Aug. 1	Aug. 10, " "
"	Atomizer	"	Aug. 10, " "
<i>Gloe. musarum</i>	"	July 27	Aug. 10, " "
"	Puncture	"	Aug. 10, " "
<i>C. lagenarium</i>	"	Nov. 1	Nov. 10, " "
"	Atomizer	"	Nov. 10, " "

pears on the tree. Edgerton₁₀₉ states that the above two organisms readily lose their virulence when grown for some time on artificial media. The *Gloeosporium* sp. from *Populus deltoides*, *Colletotrichum lindemuthianum* (Sacc. & Magn.) B. & C., and *Gloe. musarum* Cke.

**Gloeosporium piperatum* reisolated from sweet pea seedlings infected by puncture inoculation.

& Mass. failed to infect the sweet pea after repeated trials both by puncture and atomizer inoculations. In comparing Tables I and II we see that the organisms which infect the sweet pea also infect the apple, with the exception, however, of *Gloe. gossypii*, which readily infects the sweet pea by atomizer inoculation, but always fails to infect apples on the tree.




Fig. 40. Bitter rot of apple induced by the same fungus which causes anthracnose of the sweet pea, viz.—*Glomerella rufomaculans*.

The data in Table IV show that none of the organisms used could infect the Rubicon apple on the tree when the fruits were about the size of a large grape. Later, however, by June 26, the first positive infection was obtained with *Glom. rufomaculans* from the apple. At this same date all the other organisms used failed to infect. On July 15 the same condition prevailed. By August 19, typical bitter rot infections were obtained with *Glom. rufomaculans* from the apple and sweet pea, *Gloe. officinale*, *Gloe. gallarum*, *Gloeosporium* sp. from May apple fruit, and *Gloe. piperatum*. Negative results were obtained with *Glom. rufomaculans* from fig, *Glom. gossypii*, *Gloeosporium* sp. from *Populus deltoides*, *Colletotrichum lindemuthianum* and *Gloe.*

musarum. Further inoculations were made on the Rubicon apple until September when the fruits were ripe and the results were the same as mentioned above.

Field inoculations were tried on the Paradise Sweet apple with the organisms mentioned in Table IV and with like intervals of time between them. The results obtained were the same as with the Rubicon except that positive infection with *Glom. rufomaculans* from apple was not obtained before Sept. 7. Inasmuch as the Paradise Sweet is a late variety this indicates the greater resistance of late over early varieties.

TABLE IV

Species of fungus	No. fruits inoculated	Dates of inoculation	Results from inoculation
12 species named below	6 fruits for each organism	June 2	June 12, all healthy
12 species named below	"	June 6	June 21, all healthy
<i>Glomerella rufomaculans</i> from apple.....	8	June 14	June 26, all typical bitter rot
11 other species named below.....	8	"	" all healthy
<i>Glom. rufomaculans</i> from apple.....	8	June 27	June 15, all typical bitter rot
<i>Glom. rufomaculans</i> from sweet pea.....	10	"	" 8 fruits show small spots
<i>Gloeosporium</i> from May-apple.....	10	"	" 2 fruits are healthy same as above
9 other species named below.....	10 fruits for each organism	"	" all healthy
<i>Glom. rufomaculans</i> from apple.....	8	July 7	Aug. 2, all typical bitter rot
<i>Glom. rufomaculans</i> from sweet pea.....	10	"	" " "
<i>Gloeosporium officinale</i>	10	"	" " "
<i>Gloe. gallarum</i>	10	"	" " "
<i>Gloeosporium</i> sp. from May apple.....	10	"	" " "
<i>Colletotrichum phomoides</i>	10	"	" all very small spots, but typical bitter rot
<i>Gloe. piperatum</i>	10	"	" very small spots but not typical bitter rot
<i>C. gloeosporioides</i>	10	"	" same as above
<i>Glom. gossypii</i>	10	"	" all healthy
<i>Gloeosporium</i> sp. from <i>Populus deltoides</i>	10	"	" " "
<i>C. lindemuthianum</i>	10	"	" " "
<i>C. lagenarium</i>	10	"	" " "

Field inoculations were also carried on with Kieffer pear fruits. The latter being a variety which ripens very late, positive infection on the fruit with six of the organisms could not be obtained earlier than Oct. 6. The above fungi which produce the anthracnose disease of the sweet pea and the bitter rot on the apple also produce typical bitter rot on the Kieffer pear.

The foregoing experiments have conclusively proved that the anthracnose disease of the sweet pea is identical with *Glom. rufomaculans* which produces the bitter rot of the apple. Moreover, the six organisms above mentioned, which were previously considered as distinct species, are now shown through the above experiments to be identical with *Glom. rufomaculans*, since they readily produce the typical anthracnose disease on the sweet pea and the bitter rot of the apple and the pear fruits on the tree. The experiments further indicate the saprophytic nature of *Glom. rufomaculans*_{1,10} since no infection could be obtained on very young apples or Kieffer pear fruits on the tree. In the Delaware bulletin just referred to an explanation is given of the causes of the difference in resistance between different varieties of the same fruits and between young and older fruits of the same variety. If we look for an explanation as to why *Glom. gossypii* infects sweet peas and fails to infect apples and pears on the tree but readily infects the same fruits when they are picked and placed in moist chambers in the laboratory, we are brought to the following theory: It seems that the *Glom. gossypii* at one time was identical with *Glom. rufomaculans*, but that through long association with the cotton plant it has become so modified in its habits as to make it a physiological species capable of infecting the sweet pea and possibly other hosts, but having lost the power to infect the apple. From this it would seem that it is the cell contents of the host which may in some cases modify the physiological habits of an organism. To refute the above statement it could not be argued that the sweet pea can be infected by all species of *Gloeosporium*. This is not the case, since experiments have proven that only the organisms which infect the apple can also infect the sweet pea, with the above exception.

The writer hopes to continue experiments along these same lines with the object of finding out whether certain other supposedly different species of the *Glomerella* type are not one and the same.

Mode of Infection and Period of Incubation. The anthracnose of the sweet pea is mainly a disease of the tender parts of the

plant. Infection starts at the tips and the fungus works downwards invading both stems and leaflets until it reaches a node on the older parts of the stem, where it is stopped in its course. It is not infrequent, however, to find whole branches dying, and sometimes the entire plant is involved. In such cases it has been found that the plant is suffering from insect attacks, either by plant aphids (*Aphis* sp.), or more especially the red spider, (*Tetranychus bimaculatus* Haw.). These help the fungus both by weakening the host plant and by distributing the spores over its surface. The spores when germinating have no difficulty in penetrating the oldest parts of the host if it has been punctured by these insects. This explains why the plants suffer most during the hot dry weather, since at that time the aphids are most abundant. Infection often begins with the blossoms at the junction between the flower and the peduncle, in which case the blossom shrivels. The pods, also, even those which are nearly ripe, are often seen to be badly affected. Here, too, the aphids will be found to have opened the way for the fungus. These symptoms are observed in the field when infection takes place naturally, or in the laboratory where the plants are artificially inoculated with *Glom. rufomaculans* from the sweet pea. The same mode of infection and the same symptoms are observed with the other organisms which are capable of infecting the sweet pea.

The spores of *Glom. rufomaculans* from the sweet pea usually germinate in from six to twenty-four hours, according to the amount of moisture in the atmosphere. The germ tubes enter the host by breaking through the epidermal cells of either leaf or stem. In case the spore lodges on a stomate, the germ tube grows away and avoids entrance. It may be that the gases which are given off at the stomates are toxic and prohibit the entrance of the germ tube, which often breaks through the epidermal cells as soon as the spore germinates. At other times the spore germinates by sending out a short germ tube which forms an appressorium which attaches to the epidermal wall. This appressorium is then seen to germinate, its germ tube breaking through the epidermal cell.

The period of incubation varies from three to five days according to the amount of moisture in the atmosphere. The acervuli appear within five days after wilting begins unless the weather is dry, when they may not appear until eight to ten days after infection. In the field the sweet pea anthracnose is at its height during July and Aug-

ust. This is also the time when the bitter rot of the apple makes its appearance in the orchard. It is thus easy to understand how readily the natural cross inoculation may be effected.

Morphology of the fungus. The spores and mycelium of the Gloeosporium of the sweet pea do not differ from the corresponding structures of *Glomerella rufomaculans*. Sheldon first observed endospores of the Gloeosporium of the sweet pea in pure cultures, and the writer has observed the same structures in hanging drop and plate cultures of *Glomerella rufomaculans* from the apple and the sweet pea, *G. officinale* and *G. psidii*.

The formation of endospores is as follows: In some of the mycelial threads the protoplasmic content rounds itself into one or more cells, resembling chlamydospores. At the tip end of these cells a filament grows out within the empty part of the mycelial thread and at the tip of this filament the endospore is formed in the same manner as the conidia on the conidiophores of a Gloeosporium. The endospore is broken off and pushed forward for the formation of a new one. Further studies are necessary to determine the conditions necessary for endospore formation. The spores of *G. fructigenum*, *G. Gallarum*, and the Gloeosporiums from sweet pea and May apple, all germinate in the same manner, by sending out a stout germ tube. On five different synthetic media these Gloeosporiums produce growths and fructifications of the same character. The spores of the species from guava and *Persea* germinate by sending out a very thin germ tube. They also differ in manner of growth and fructification on the synthetic media from the other organisms used. Until the perfect stages are found, it appears from these studies that we are justified in considering *G. gallarum*, *G. officinale* and the Gloeosporium from the May apple fruit as one and the same with *Glomerella rufomaculans* (Berk.) Spaul. & von Sch. of the apple and the sweet pea.

Life History. In order to determine whether the disease is carried over with the seeds of the sweet pea, a large quantity of diseased pods were collected and kept over winter, some in the laboratory and some out of doors. Spores from both lots of materials were tested Nov. 22, Dec. 22, 1910, Jan. 22, Feb. 22, March 15, 20, 24, April 20, May 21, and June 20, 1911. In all cases the spores germinated well and produced normal colonies on bean agar. Spores of cultures obtained from the sowing of June 20, 1911, readily infected healthy

sweet pea seedlings, which were grown in pots in the laboratory. This indicates that the organism is carried over the winter as viable spores on the pods and on the seeds, and there is very little doubt that the disease is introduced into new localities through diseased seeds.

MOSAIC DISEASE OF THE SWEET PEA

Brief History. The first study of the disease was made by Mayer₁₁₁ in 1866 on the mosaic of tobacco. Mayer found that the disease was not induced by insufficient mineral nutrients. He also found the disease to be distributed over large areas irrespective of soil conditions and further proved that the juice of the leaves of affected plants, when injected into healthy leaves would reproduce the disease in from ten to eleven days. Mayer found that a temperature of 60° C. does not destroy the infectious nature of the juice, but that a temperature of 80° kills it. Mayer could not find any animal or fungus parasite to be associated with the disease, altho he believed the true cause to be bacteria which could not be isolated.

In 1892 Iwanowsky₁₁₂ confirmed Mayer's work. He too was unable to isolate the specific germ on artificial media but states that he saw the bacteria and proved their presence in the tissues of the affected host. In 1894 Prillieux and Delacroix₁₁₃ found this same disease on tobacco in France.

In 1897 Marchal₁₁₄ described under the name of "La mosaïque du tabac" a disease similar in appearance to that described by Mayer₁₁₅. Marchal claims to have obtained from diseased leaves a motile bacillus forming chains in culture media, and capable of reproducing the disease by inoculating with a pure culture.

In 1898 Beijerinck added considerable to our knowledge of this disease. He apparently proved the absence of bacteria in the development of the disease. When the juice of diseased plants is passed through filters, the liquid, while remaining perfectly clear and free from bacteria still retains the power of infection. He found that only growing meristematic tissue could become diseased. Among many other things he also found that soil from diseased plants may infect healthy plants. He further showed that the infective material could be transported through considerable distance without losing its virulence. He assumed the virus to be a non-corpuseular fluid like material which has the power of growth when in contact in a sort of sym-

biosis with growing cells. He called it therefore "a living fluid contagium."

In 1902 Woods₁₁₆ concludes that the disease is due to the presence of oxidizing enzymes in the plant; these he designated as oxidase and peroxidase.

In 1904 Selby₁₁₇ refers to the mosaic disease of tobacco as a non-parasitic disease, he, accepting the conclusions of Sturgis and Woods.

In a recent paper, Allard₁₁₈ has succeeded to transfer by means of inoculation the mosaic of tobacco to other solanaceous plants of the following genera: *Nicotiana*, *Lycopersicum*, *Petunia*, *Physalis*, *Datura*, *Hyoscyamus*, *Solanum* and *Capsicum*. T. Allard also found aphids to be carriers of the disease.

Mosaic Disease of the Sweet Pea. As far as known there is no mention made before of the mosaic disease of the sweet pea. The writer first made a study of the disease in the summer of 1912 and his first published report appeared in the *Florist Exchange*₁₁₉.

Like the anthracnose, the mosaic of the sweet pea is a very important disease. Both out door and greenhouse plants are alike subject to it.

Symptoms. Mosaic is readily distinguishable by a yellow dotting or mottling of the leaf, presenting in some instances a beautiful mosaic structure, hence its name (Fig. 41). Affected leaves seem to linger for a time but they eventually lose all their chlorophyll and soon drop off. Another symptom of this disease is a curling of leaves (Fig. 42), resembling the curling induced by the green aphids, but in this case the aphids had no association with it. The disease makes its appearance after the seedlings are from three to four weeks old. Often, the disease is so bad and the curling so pronounced that the plants thus affected cannot make any headway and remain dwarfed. An attempt is made by these curled parts to produce a few flowers, but the later are borne on very short peduncles as compared with the long peduncles of healthy plants of the same variety. Frequently, however, the affected plants outgrow the disease entirely, and thus a distinct line of demarcation is observed between the previously diseased part and the healthy part of the new growth. At other times infected plants keep on growing and even flowering, with the disease keeping pace.




Fig. 41. Sweet pea mosaic disease. (a) (b) leaves showing mosaic; (c) healthy.

Pathogenicity. Like peach yellows and mosaic disease of tobacco and tomato, this disease of the sweet pea, too, can be reproduced by a puncture with a sterile needle from the diseased leaf into

a healthy leaf. No organism could be obtained in culture, nor could it be detected with the microscope. Nevertheless, this disease is contagious, as is the peach yellows. When the disease first made its appearance in our experimental sweet pea field, the diseased areas



Fig. 41 Mosaic disease causing dwarfing of the plant and rolling of the leaves.

were immediately located in order to learn something of its spread. They formed two small areas, one in about the center of the field, the other in the southeast corner. Within ten days another survey was made and the whole field was found to be infected. With the exception of the dwarf Cupid varieties, which are seemingly immune, all the rest were found to be affected with the mosaic. When first investigating this disease I thought that, perhaps, this mottling of the leaves was merely a variegated condition. We also thought that perhaps the curling of the tender tips as well as the mosaic effect was due primarily to the presence of aphids, which at the beginning of the season were so plentiful. Experiments were then undertaken to determine definitely these points. Accordingly, sterile pots with sterile soil were isolated in a glass chamber and the plants were allowed to grow for three weeks to see if any disease would develop on them. However, these plants remained very free from any disease. The pots with plants were then divided into four lots; into Lot I were introduced a few stem mothers of aphids from affected mosaic plants in the field. In Lot II were introduced a few stem mother aphids from apparently healthy plants in the field. The plants in Lot III were punctured with sterile needles and by pricking a mosaic affected leaf, and then puncturing with the same needle the healthy leaves. Lot IV was punctured merely with the sterile needle, the plants of this lot were designed to serve as checks. In each lot there were two pots with plants in order to duplicate each experiment. After ten days the lot which were inoculated with the aphids from the diseased and healthy plants both began to show the symptoms of mosaic. This, therefore, would appear to show that the mere puncture of aphids would be responsible for the mosaic disease. However, this is not the case, as we will soon see. Moreover, it is easy to suppose, and that on very good ground, that the aphids taken from seemingly healthy plants in an infected field might themselves have been infected before. But this would be no valid proof. Lot III, which was infected with needle punctures from diseased leaves, began after ten days to show the mosaic disease, while the check punctures remained all healthy to the end of the experiment. This definitely proves that the aphids are not the cause of the trouble but are merely the carriers of the mosaic disease. It seems, therefore, that any steps taken to control the aphids may also serve to control the mosaic. From this, too, it is evident

that not only the aphids, but also any biting or sucking insect may help spread the disease.

It has been definitely proven that the mosaic disease is contagious, since it can be produced at will by artificial inoculations. The symptoms produced in artificially inoculated plants are similar to those in the field, namely, a yellowish spotting or mottling of the leaves and a tendency of the leaves of the tips of the plant to curl.

Cause of the Mosaic Disease. Mention has been made of previous workers who with ourselves have definitely proven the infectious nature of the disease. Attempts to prove the cause of the trouble have resulted in failure. Mayer¹²⁰ failed to find the association of animal or fungus parasites with the disease. He thought that bacteria were the cause of the disease, but all inoculations with bacteria cultivated from the surface of diseased leaves, and with mixtures of different bacteria gave negative results.

Beijerinck¹²¹ disproved the theory that the cause of the trouble was bacteria, by showing that the juice of diseased plants filtered through Chamberland filters while remaining perfectly clear and free from bacteria still retained the power of infection.

Sturgis¹²² in his conclusions states as follows:

“It (mosaic) is not caused by predaceous insects, nematodes, or parasitic fungi.

“Bacteria have not been associated with the disease but no critical method for their isolation or culture has been applied, and therefore the question of their influence cannot at present be answered.

Woods¹²³ believes the disease to be of a physiological nature and of enzymic activity. Woods claims to have reproduced the disease several times by merely removing the tip of a rapidly growing plant.

Suzuki¹²⁴ in his studies of the so-called mulberry dwarf troubles in Japan (all evidence seems to show that this disease is of a similar nature as our mosaic), concludes that the principal cause of the disease is due to the practice of subjecting the mulberry trees to repeated low cuttings, thus removing the reserved food which is required for growth. Woods believes that it was the enzyme (peroxidase) of the leaves that induced the disease because he claims that he induced the disease artificially by injecting into a healthy plant the juice from another healthy plant. Woods found more peroxidase in diseased than in healthy plants. “It is through the introduction of the enzyme

into the infected host that pathological changes are brought about which result in an increase of the normal enzymes of the cell and the decrease of available reserve food. When this condition is reached, it is very difficult for the plant to outgrow the trouble." Accepting this hypothesis, Woods is at a loss to explain how the disease is spread.

In my investigations repeated trials failed to reveal the presence of either fungus or bacteria in culture. Nevertheless I do not believe with Woods that the disease is physiological and enzymic. I strongly believe the trouble to be either bacterial or protozoic* and the pathogenic nature of the disease strongly points to this conclusion. That all attempts to obtain a living micro-organism in pure culture have failed does not argue against the possibility of its being either bacteria or protozoa, but simply that our present cultural or filtering methods are not suitable for its detection or retention. Previous to the discovery of *Bacterium tumefaciens*, by E. F. Smith, no one suspected the crown gall of plants to be of bacterial origin, cultural attempts in each case failed to reveal the organism.

As I have already indicated, Beijerinck showed that the juice of diseased plants when filtered through Chamberland filters, while remaining clear and free from bacteria still retains the power of infection. This proof too does not argue against the possibility of the bacterial or protozoic nature of the mosaic, because the former may be even smaller than bacteria and readily pass through the Chamberland filters. It is therefore possible that Beijerinck's filtered fluid was contaminated with the pathogen and this is why the filtered fluid retained its pathogenic nature. Neither can we accept Wood's statement that healthy plants when cut back develop the disease unless we admit again of the possibility of contamination. If the mosaic pathogen is not present or has not made its appearance on a certain host, the latter can be cut back time and again without the disease ever making its appearance. Before I introduced the disease through puncture inoculation with diseased tissue in the laboratory, I cut back my experimental sweet pea plants in order to prevent them from growing too high and altho I have practiced this very often I have never had a case of mosaic develop from this operation. It is also difficult to believe with Woods that the disease is of a physiological and enzymic nature. Enzymes in plants are natural factors of im-

*The idea of the protozoic nature of mosaic was suggested to me by Dr. T. F. Manns.

munity to protect that plant from disease. Diseased plants will naturally possess a higher enzymic content just as much as the leucocytes in the human blood increase in case of wounds, etc. That this peroxidase of Woods from diseased plants should possess the power of infection is very possible as I have already pointed out the former may be contaminated. That the peroxidase of healthy plants when inoculated into a healthy plant should be able to produce the disease as Woods claims to have done, is probable, only if we admit of possibilities of contaminations which may have been the case. As already stated, the pathogenic nature of the disease points to a living organism as being the cause.

In Wood's enzymic explanation we have no means of accounting for the spread of the disease. Under pathogenicity we have shown that green aphids carry and spread the disease from leaf to leaf and from plant to plant. This by itself is sufficient proof that a living organism is the cause of the mosaic, for if as Wood maintains, the peroxidase of a healthy plant when inoculated into a healthy host will reproduce the disease and if there be no chemical difference between peroxidase of healthy and diseased plants, then why is it when green aphids from perfectly healthy plants are transferred to healthy sweet peas the disease never develops but the mosaic readily makes its appearance by introducing green aphids from diseased plants to healthy ones. If the aphids can carry the peroxidase from diseased plants why do they not carry the same from the healthy plants while sucking their juice? This to my mind is the strongest argument against Wood's physiological and enzymic nature of the mosaic, and on the other hand it strongly points to the probable activity of a living bacterial or protozoic organism. Aphids are not the only agents capable of carrying and distributing the disease, for there are others which may do it. Among the biting insects we have the "corn root worm beetle" (*Diabrotica longicornis*), the striped potato beetle (*Epicanta vittata*) and several others which feed on the sweet pea and at the same time help to distribute the mosaic.

Transmission of the Mosaic through Seed or Soil. Beijerinck claims that soil around diseased plants may infect the roots of healthy plants. In order to determine whether the disease is carried in the soil, the following experiments were tried. A number of sterile pots were taken and arranged in groups. Group A. consisted of pots filled with sterile soil, on the level surface of which leaves infected with

mosaic were spread. Sterilized seeds were sown on the surface of the leaves and the infected leaves and seeds were covered with a 2-inch layer of sterile soil.

Group B. consisted of 2 pots filled with sterile soil but the latter was mixed thoroughly with a quantity of mosaic infected leaves. These pots were sown with sterilized seeds and covered with a 2-inch layer of sterile soil.

Group C. consisted of two pots filled with sterile soil and sown with sterile seeds and then thoroughly watered with water from a vase in which mosaic infected plants stood for two days.

Group D. consisted of two pots filled with soil taken from the field and from a spot where mosaic infected plants grew.

Group E. were untreated checks, that is, two pots with sterile soil were sown with sterile seeds and placed at a distance from the infected series. All the pots were then watered with distilled water up to the end of the experiment. The seeds in all the groups germinated and the plants were allowed to grow for ten weeks when they had attained considerable size and in no case did the mosaic appear, thus apparently proving that the mosaic soil is not a factor in carrying the disease. There is no evidence either that the mosaic is carried with the seeds for in no case did the disease appear in the laboratory where unsterilized seeds were used.

Diseases of the Sweet Pea not known to be present in this Country

In an article in "The Sweet Pea Annual," Massee¹²⁵ describes the following diseases, which as far as known, have not as yet made their appearance in this country on the sweet pea.

Pea Blight (*Peronospora trifoliorum*)

Pea Spot (*Ascochyta pisi*)

PEA BLIGHT, *Peronospora trifoliorum* De By.

According to Massee, this is the most destructive disease to peas, lupines, and to most of the pea family. The disease may appear and spread quickly when the plants are only a few inches high, or it may attack older plants. In dry weather the mycelium of the fungus present in the tissue spreads throughout the leaf,

which soon assumes a sickly yellow-green color, and finally bleaches, shrivels and dies without showing any, or only a small amount of mould on the surface. In damp, dull weather infected leaves show yellow patches, which soon becomes covered on one or both surfaces with a very delicate greyish-lilac mould. The summer spores are produced on the leaves, or on any other part of the host. The winter, or resting spores, are imbedded in the tissue of the host that has been previously killed by the fungus. The resting spores have a very thick smooth brown wall. *Peronospora viciae* is also stated to be able to produce a disease on sweet peas.

PEA SPOT, *Ascochyta pisi* Lib.

According to Masee, this disease also attacks the French beans and several other leguminous crops. The first indications of disease on the pods is the appearance of pale green spots of variable size and irregular shape. These blotches continue to increase in size for some time and eventually become whitish, bordered with a dark line, and have the surface studded with minute black points which are the pycnidia of the fungus.

C. Diseased Seeds

Under the discussion of the anthracnose disease I have shown that the fungus (*Glomerella rufomaculans*) is transmitted with the seed. In that case infection starts at the pods and the fungus works inwards gradually penetrating the seed coat and the seed proper (Fig. 38). Such seeds when harvested have a shriveled appearance and when planted with healthy seeds introduce the fungus in the soil and then the disease begins to attack the adjacent seedlings, thus spreading throughout the whole field.

Another disease that may be transmitted with the seeds is the "Streak". In examining infected plants we can readily see the organism (*Bacillus lathyri*) invade the pods and then work into the seeds. Pure cultures of the organism may readily be obtained by surface sterilizing an infected pod, picking out the seed with sterile forceps and then dropping the same into a plate of media. We have as yet no data to prove that the organism can survive the drying when it is only present on the surface of the seed coat. However, as is often the case, the organism works through the seed coat and into the seed. Under such conditions it is very probable that it may be carried over

winter while present in the seed tissue. Such seeds when planted introduce the parasite into the soil and from there the disease gets a foothold to carry on its destructive work.

Sweet pea seeds as we buy them from the seedsmen are put in small paper packages. During my work on the sweet pea diseases I had occasion to open two thousand of such packages. In very few




Fig. 43. Fusarium and Botrytis fungi from shriveled and non-germinated sweet pea seeds.

cases were all the seeds plump and full. A certain percent were shriveled and gave the appearance of being diseased. It was thought, therefore, advisable to make a study of such shriveled seeds to deter-

mine whether or not they are disease carriers and the nature of the pathogens. The technique employed was as follows:

Fresh packages of sweet peas were opened and the percent. of shriveled seeds in each package determined, labeled and separated. The next step was to soak each lot of seed with a 5% formaldehyde solution for $\frac{1}{2}$ hr. and then to wash it three times in sterile water in order to remove all trace of formaldehyde. Then with a sterile forcep, each lot of seeds thus treated was dropped into a petri dish containing nutrient agar that had been melted and cooled to the proper temperature. As the agar solidified the plates were placed in the incubator and kept there for one week. Observations were made every other day to determine the percent. of germination and to make transfers into slant tubes of agar of all fungus or bacterial growth which appeared on the shriveled seeds (Fig. 43). The results obtained are given in Table V from which will be seen that a large percentage of

TABLE V

Name of variety in package	Per cent of shriveled seeds in package	Per cent of germination of shriveled seeds	Kind of fungi obtained from non-germinated shriveled seeds
King Edward VII	13%	33%	None
Gray Friar	1/100%	0	"
Aurora	9%	20%	Botrytis cinerea
Apple Blossom	12%	6%	None
Emely Henderson	18%	100%	"
Henry Eckfort	15%	4%	Fusarium sp.
Jeannie Gordon	5%	100%	None
Dorothy Eckfort	13%	80%	Fusarium sp
Hellen Pierce	10%	80%	" "
Cocceinae	6%	90%	" "
Katherine Tracy	15%	40%	Alternaria
George Herbert	19%	0	Bacteria
Black Knight	4%	100%	None
Jeannett Scott	21%	100%	"
Dobbies Mid Blue	3%	100%	"
Blanche Burpee	12%	100%	"
America	9%	100%	"
Blanche Ferry	1%	100%	"
Mrs. A. Watkins	7%	60%	Fusarium sp.
Countess Spencer	15%	18%	Botrytis
Boltons Pink	1%		Clanostachys sp
Countess Cadogan	12%	9%	None
Agnes Eckfort	16%	8%	Fusarium
E. J. Castle	3%	1%	"
Mrs. Collier	10%	50%	"
Burpees Midnight	0	0	—

the so called shriveled seeds readily germinated. It was further observed that these before germination became plump and resembled in every respect the germinated healthy seeds which were treated in the same way as the shriveled and were run as checks. It seems then that shriveling in the seeds is merely correlated with a loss of water. Whether shriveled seeds in the long run produce weaker plants has not been determined. Observations so far have shown that the seedlings from the germinated shriveled seeds were in every way equal in vigor to the seedlings of the germinated plump seeds.

Of the non-germinated shriveled seeds, those which remained free from fungous growth can be classed as hard seeds and those probably would have germinated if the seed coat had been pierced. The germinated seeds which showed growth of fungus or bacteria were at once seen to become soft and rotted. Of the organisms isolated from the non-germinated seeds there were two species of *Fusaria*, one species of *Alternaria*, one species of *Clonostachys*, *Rhizopus nigricans* and *Botrytis cinerea*. These after repeated trials failed to infect healthy sweet pea seedlings, thus seemingly proving that they are saprophytes and of secondary nature. Their presence on some of the weak and non-germinated seeds, no doubt helped in the decay of the former, but they fail to play the role of active parasites on growing plants.

II. Bacterial Diseases

So far only one kind of bacterial disease of the sweet pea has been observed and that is the "streak" disease. This has been determined to be caused by a newly described organism, viz. *Bacillus lathyri* Manns and Taubenhau.

STREAK IN ENGLAND

Historical. In correspondence with Mr. T. A. Weston of Orpington, England, the former states that the disease was first observed by H. J. Digges of Dublin in about 1904 or 1905.

In 1906 T. A. Weston₁₂₆ gave the name of "streak" to the disease referred to above (the cause of the disease was not given). In 1908 Masee₁₂₇ in a letter to a correspondent who had sent in diseased specimens replied, "the disease is of a physiological nature" and "brought about by over feeding." In 1912 Chittenden₁₂₈ believed that the fungus *Thielavia basicola* was the cause of the "streak". For the first two years of his inoculation experiments Chittenden

failed to infect healthy seedlings with the *Thielavia* fungus. It was only by overwatering his plants that he succeeded in getting some root infection, as he states: "To sum up, as far as our experiments go, the 'streak' disease is brought about by the attack of the fungus *Thielavia basicola* on plants that have received some check at the root." Chittenden has failed to reproduce the typical "streak" but merely the *Thielavia* root rot as I have already indicated under my description of the above fungus. In the same year Massee₁₂₉ again studied the disease and attributed it to *Thielavia basicola*.

In 1912 Dyke₁₃₀ found *Macrosporium solani* constantly associated with the disease, and believed it to be the cause of the trouble.

STREAK IN THE UNITED STATES

As far as I know, up to 1913 there were no American references to this disease.

Massee's₁₃₁ short note led me to believe that the disease was physiological. Under Mosaic disease, I have shown that some workers believed it to be a physiological disease. While working on the mosaic, and having the "streak" in mind, I₁₃₂ made the following statement:

"In England the sweet peas suffer from a disease known as 'streak'. This disease is very much dreaded by English gardeners, as it causes great losses. From the description given of that disease it seems to be similar to the new mosaic disease of this country. However, we refrain from passing final judgement until we have the opportunity of seeing the English specimens and of making comparisons. In England the streak disease is attributed to a fungus *Thielavia basicola*, which attacks the roots. In our investigations we have not found the *Thielavia* fungus or any other organism associated on the roots of mosaic affected plants. In fact, such affected plants were found to have as normal a root with as much in the way of legume nodules, as the healthy ones. If our mosaic disease proves to be the same as the streak disease of England, it will be safe to assume that the *Thielavia* in England is secondary and merely follows the already weakened mosaic affected plant."

The above was published on July 20, 1912, and the statement was made before the American Sweet Pea Society at Boston early in July of the same year.

In the middle of July of 1912, I first noticed a peculiar disease on the stems of sweet peas grown in our experimental field. The disease

was characterized by dark streaks running along the stems. This suggested at once the possibility of its being the "Streak" disease of England. Diseased specimens were brought in the laboratory, and microscopical crush mounts were made, but no organism was found that would give any clue as to the nature of the trouble.

In submitting specimens to Prof. Manns and in consultation with him, it was suspected that the trouble might be of bacterial origin. We at once started to make cultures from the diseased material and a pure culture of a yellow bacterium was obtained. Platings made from diseased material from widely separated gardens gave a pure culture of the same bacterial organism. Cultures made from diseased material sent in by Wm. Sims of Boston, Mass. and by T. A. Weston of Orpington, England, all gave an organism similar to that isolated from the home material. Stains made of this organism revealed a bacillus sp. In the meantime Prof. Manns recalled that he had seen a similar streak disease on clovers in Ohio. A search for plants infected in this way showed it upon clovers and upon some other leguminous plants in the vicinity of Newark, Del.

Inoculations made with the bacterial organism on the sweet pea reproduced the typical "streak" disease. That the "streak" is present and widespread in the United States there is no doubt. I have seen it in widely different localities in Delaware, in Pennsylvania, in Massachusetts, and in New York. A letter addressed to me by C. C. Morse & Co., of California, gives the following information:

"The 'streak' disease has not appeared in California as yet, and this is accounted for by the fact that growers so far have not gotten into methods of over-manuring their grounds."

As previously stated, the American literature contains no reference to the disease. Manns and Taubenhaus* published their first account of the disease in the *Gardeners' Chronicle* of England, announcing the "streak" as being a bacterial disease and the parasite a newly described bacillus, giving it the name of *Bacillus lathyri*.

Symptoms.† Like the Bacteriosis of beans, streak makes its appearance in the season of heavy dew. On the sweet pea the disease usually appears just as the plants begin to blossom; it is manifested by

*Manns, T. F., and Taubenhaus, J. J. "Streak, a Bacterial Disease of the Sweet Pea and Clovers." *The Gard. Chron.*, London, Apr. 5, 1913.

†Abstracted from above article.

light reddish-brown to dark brown spots and streaks (the older almost purple) along the stems, having their origin usually near the ground, indicating distribution by spattering rain and infection through the stomata. The disease becomes quickly distributed over the more mature stems until the cambium and deeper tissues are destroyed in continuous areas, when the plant prematurely dies. From the stems the disease spreads to the petiole, flower, peduncles and pods. The symptoms in these cases being similar to those on the stems. On the leaves, however, the disease appears as small roundish spots which gradually coalesce, and eventually involve the entire leaf, which when killed presents a brownish dark appearance."

Pathogenicity. The pathogenicity of the causative organism may be proven by diluting a pure culture of the organism in sterilized water and by spraying on the plants with an atomizer. This should be done in the evening when the temperature is cooler and there is less chance for evaporation of the applied infectious liquid.

The disease makes its appearance from seven to ten days after artificial infection and the symptoms are similar to those produced in nature. The organism may be reisolated from the artificially infected plants and the disease induced again at will on healthy plants, in each case the check remaining healthy.

Natural or artificial infection can only take place on mature plants which have started to bloom. All attempts to inoculate plants in all stages of growth previous to the blooming has failed. It seems that the host possesses certain protective properties previous to the blossoming which inhibits the growth of the parasite. Failure to infect young plants was not due to abnormal conditions or to bad technique. The disease in the field does not make its appearance until the plants have started to blossom.

Isolation and Morphological Studies. Over 1,500 plate cultures of beginning or young lesions were made from the several hosts. The organisms may almost invariably be taken in abundance in pure cultures from the beginning lesions in the stems of sweet peas when the surface is properly sterilized.

The isolation work readily indicated the parasite to be a bacterium; a yellow organism which grows luxuriantly upon all the nutrient media, and especially rapid upon nutrient media containing sugars.

On standard nutrient glucose agar the colonies appear within 24 to 36 hours. The center becomes granular and the colonies have a marked tendency to become stellate or auriculate.

Morphological studies show the organisms to be a comparatively small rod-shaped bacillus, which in fresh cultures is rarely found in chains, and seldom united in twos or fours. The flagella are not easily demonstrated; they are shed so readily that usually not more than two to five may be stained, and these are generally quite short. However, when proper material is selected, carefully fixed and stained, the flagella may be demonstrated to be very long and delicate, and to number 8 to 12, well distributed peritrichially*.

III. Physiological Diseases

By physiological diseases of the sweet pea we mean all disturbances in the plants which are not induced by fungi, bacteria, insects, or any other parasites but is expressed in a disturbance in the metabolism of the plant. This disturbance seldom results in the sudden death of the plant.

Under Physiological diseases I will consider the following two troubles: 1. Bud drop. 2. Arrested development.

BUD DROP

As the name implies the young flower buds at a very early age turn yellow and drop off. This drop should not be confused with the drop produced by the anthracnose disease. In the latter case, the flower develops into a normal spike but it is attacked soon by the fungus *Glomerella rufomaculans* which girdles it at a point of attachment between the flower and the peduncle. In this case the flower often drops off leaving behind the beheaded peduncle. In the latter case, however, the minute young flower bud never develops, instead it turns yellow and drops off.

There seems no doubt that the drop is a physiological disease and is induced by an unbalanced condition of food elements in the soil. This may occur in a soil that has been excessively fed or in a soil that is lacking in plant food.

The following extract of a letter from a grower whose plants have suffered severely from the drop and who gives the history of his soil

*A more detailed account of this organism will soon appear in a Delaware Bulletin by Dr. T. F. Manns.

treatment will help to confirm the physiological nature of the disease.

“The soil is rich and is located in a valley near a creek, the sub-soil is similarly rich but is not of a mucky nature. Before sowing the peas the field was trenched and a thick layer of pig manure mixed with a little hen manure was put at the bottom of the trench. The manure was mixed up with soil in the trench and the seeds sown thereon. After germination and when the plants reached from 8 to 10 inches, a commercial fertilizer (kind of fertilizer not stated) was worked in at both sides of the row. A short time before blooming, a layer of coarse stable manure was put around the plants to serve as a mulch. During blossoming time the flower buds began to drop off heavily and what promised to be a successful crop of blooms looked as though it would result in total failure.” It is here very evident that the plants were supplied with too much nitrogenous matter but with little of potash and other mineral elements.

Cultures made from these fallen buds failed to produce an organism of any kind.

In order to remedy this trouble Prof. T. F. Manns suggested the application of 150 lbs. muriate of potash and 600 lbs. acid phosphate per acre. This treatment was followed out by the grower and the drop ceased within a week resulting in a perfect crop of flowers.

On poor soils I have often seen this same “flower drop” and it is also especially evident where sweet peas are grown in pots and in poor, light, gravelly soil in the laboratory. An application of a balanced fertilizer to these pots readily helped the plant to overcome the bud drop.

ARRESTED DEVELOPMENT

This trouble, too, is a physiological disease and is induced by the use of excessive fertilizers. The following facts from the letter of a grower who has suffered from this trouble will also help to confirm the belief in the physiological nature of the disease.

The seeds were sown Nov. 1st in pots and planted Dec. 15 in the beds in the greenhouse. Previous to the planting the beds were well manured with horse manure which was applied six months before planting. Besides this, wood ashes were also applied to the beds at the rate of 1500 lbs. to 4500 sq. feet of bed space. This would be equivalent to nearly seven and one-half tons per acre. About one

month after planting some of the plants turned yellow and died. Upon examining the dead diseased specimens the plants were found to be dwarfed with a sickly yellowish appearance. The roots presented a burned appearance suggesting the attacks of *Thielavia*. Microscopical examinations and cultures made from the diseased tissue did not reveal the presence of any parasite which could be associated with the soil. In submitting some of the soil to Prof. T. F. Manns for examination, he found it strongly alkaline. Hard wood ashes contain about 30% caustic lime and from 5 to 12% potash. According to Prof. Manns it was the excess of these elements in the soil that made it so highly alkaline, and this condition injuriously affected the plants. This kind of injury could be considered purely physical since it is brought about by the exposed surface of the roots to an alkaline substance. Nevertheless any injury which interferes with the metabolism of the roots is reflected in a derangement of the metabolism of the plant. The resulting injury is therefore of a physical nature.

As a remedy for this trouble Prof. Manns advises the use of acid phosphate, followed by a good drenching of water. This will neutralize the alkaline effect of the soil and also help to balance the plant ration.

METHODS OF CONTROL

Under methods of control the following lines of investigation have been carried on:

Resistant varieties

Seed treatment

Treatment of soil with chemicals

Studies of the fungicidal value of some chemical poisons

Formaldehyde treatment of soil

Steam treatment of soil

RESISTANT VARIETIES

To test out the resistance of different varieties of the same host to a certain parasite, the general practice is to plant in the field the varieties to be tested and to allow full sway to the natural causes of

infection. At the end of the growing season an estimate is taken of the per cent of infection of each variety and on that basis a scale of resistance is formulated. While this method by itself is fairly valuable, the method nevertheless is unreliable because a certain variety which under the above test proves highly immune or highly susceptible to disease will, under different conditions of climate, etc. prove the opposite of what it has promised to be in its first trial. The reason is very obvious. No two varieties are alike as well as no two individuals are alike. Conditions in the field are not always ideal for every variety of a certain host to become susceptible to disease. If this were the case we would have all our crops ravaged by pests. In order to make a reliable test of the resistance of different varieties it is necessary to have the same conditions of soil and care and then to submit the varieties to the severest test by making all conditions ideal for the parasite to attack the plants. To carry out this idea, I planted 100 sweet pea seeds of each variety to be tested in sterile soil and pots in the laboratory. Previous to the planting, the seeds were sterilized by being soaked in a 5% formaldehyde for $\frac{1}{2}$ hr. The seeds of the different varieties did not all germinate evenly due to the hardness of the seed coats in some seeds but eventually they all germinated. When the plants were eight weeks old each pot with its different variety was well watered and then covered with a bell jar. The latter was sterilized by being previously washed with a 1-1000 mercuric bi-chloride solution and then rinsed with distilled water. The plants remained under the bell jars for 48 hours, where all were seen to be uniformly covered with drops of water. A large amount of moisture accumulated under the bell jars and this was plainly visible by the drops of water standing on their walls. Under such conditions of moisture as described above, infection readily takes place. The infecting material chosen for this purpose was the fungus *Glomerella rufomaculans*, which causes the anthracnose disease. Accordingly a heavy suspension of spores from a pure culture was diluted in sterilized water and then applied to the plants by means of an atomizer. The inoculated plants were covered again for 48 hours. After that the bell jars were removed and the plants left uncovered. The result of the experiment is given in Table VI.

From Table VI it is seen that of all of the varieties tested not one of them proved to be entirely resistant. On the other hand it is seen

that the percent of infection differs with the variety. This means then that in each variety there are certain individual plants which are resistant to this particular fungus tested. Attempts were made to reinfect those plants which remained healthy after the first inoculation and it was found that nearly 50% in each case was infected and the rest remained resistant and continued to grow well. This conclusively shows that while no one variety is entirely immune to a disease yet there are nevertheless certain individuals of that variety which have developed the power of resistance. It is well known that if a plant is resistant to a particular disease, it may be very susceptible to another disease. The problem, therefore, is to test the desired va-

TABLE VI

Name of variety (seedlings six weeks old)	Fungus used (spores in suspension of water)	Per cent of infection
King Edward VII	<i>Glomerella rufomaculans</i>	90%
Gray Friar	" "	80%
Aurora	" "	93%
Apple Blossom	" "	90%
Emely Henderson	" "	40%
Henry Eckfort	" "	100%
Jeannie Gordon	" "	98%
Dorothy Eckfort	" "	100%
Hellen Pierce	" "	92%
Coccineae	" "	70%
Katherine Tracy	" "	70%
George Herbert	" "	60%
Black Knight	" "	80%
Jeanett Scott	" "	72%
Dobbie Mid Blue	" "	40%
Blanche Burpee	" "	90%
America	" "	100%
Blanche Ferry	" "	100%
Mrs. A. Watkins	" "	100%
Countess Spencer	" "	90%
Black Michael	" "	60%
Bolton's pink	" "	80%
Countess Cadogan	" "	100%
Mrs. Bieberstedt	" "	20%
Agnes Eckfort	" "	100%
Burpees Dainty	" "	100%
E. J. Castle	" "	1%
Glady's Union	" "	4%
Captain of the Blues	" "	60%
Duke of Westminster	" "	80%
Mrs. Collier	" "	70%
Burpees Midnight	" "	40%

rieties with all the known diseases to which they are subject. In these tests all the immune individuals must be selected and by crossing and

selecting we will build up a strain which will possess complete immunity to all or at least to most of the diseases to which that variety is subject.

SEED TREATMENT

I have previously shown that the sweet pea seeds are capable of carrying and of introducing one of the most dreaded diseases of the sweet pea, namely, the anthracnose (*Glomerella rufomaculans*). Sweet pea seeds were also found to carry several species of fungi which seem to be unable to assume the role of parasites. Nevertheless, the time may come when these fungi may become parasites of the sweet pea.

It was, therefore, thought necessary to devise some means of treating the seeds which would kill all external as well as internal parasites, and at the same time not inhibit the germinative power of the same. The following are the methods which have been tried:

Effect of temperature

Effect of sulphuric acid treatment

Effect of formaldehyde treatment

Effect of Temperature on Seed. In order to test the effect of temperature the following experiments were tried. Ten different varieties were thoroughly mixed and lots of 100 seeds each were picked out and put in pieces of cheese cloth and tied up with a string. The experimental temperatures of the water used were boiling water, 90°, 80°, and 60° C. The seeds were immersed in the water with the varying temperatures and kept there for different intervals of time, as is indicated in Table VII. In each test a duplicate series was always made, i. e. using two packages of 100 seeds each. The per cent of germination in each case expresses the average taken from each duplicate series. After each treatment seeds were placed in sterilized petri dishes containing moistened filter paper which had been previously sterilized by being placed in boiling water for two minutes.

After placing the seeds in the petri dish more sterile water was added in order to secure the amount of moisture necessary for germination. The plates were then placed in the incubators for 10 days, observations being made every two days. The result of the experiment is given in Table VII.

TABLE VII

Time Heated	BOILING WATER											90° C.											80° C.					70° C.					60°		Check										
	1 Sec	2 Sec	4 Sec	8 Sec	16 Sec	32 Sec	1 Min	2 Min	5 Min	10 Min	12 Min	14 Min	16 Min	18 Min	20 Min	25 Min	3 Min	5 Min	8 Min	10 Min	15 Min	20 Min	1 Min	2 Min	5 Min	10 Min	15 Min	20 Min	1 Min	2 Min	5 Min	10 Min	15 Min	20 Min		1 Min	2 Min	5 Min	10 Min	15 Min	20 Min				
Per Cent of Seed Germination	61	60	12	12	0	0	0	0	0	0	0	0	0	0	0	0	51	43	40	18	2	1	70	54	42	20	70	61	49	82															
Nature of Growth of the Germinating Seeds	g	g	w	k	k	k	k	k	k	k	k	k	k	k	k	k	g	w	w	w	w	w	w	w	w	w	w	w	w	w	w	w	w	w	w	w	w	w	w	w	w				
Behavior of the Non-Germinated Seeds	k	k	k	k	k	k	k	k	k	k	k	k	k	k	k	k	k	k	k	k	k	k	k	k	k	k	k	k	k	k	k	k	k	k	k	k	k	k	k	k	k	k	k		
Nature of the Fungus Growth on the Non-Germinated Seeds	r	r	s	b	b	r	p	b	p	b	p	p	p	p	p	p	r	p	p	p	p	p	p	p	p	p	p	p	p	p	p	p	p	p	p	p	p	p	p	p	p	p	r		
Treatment Successful or Not	partial	partial	on	on	on	on	on	on	on	on	on	on	on	on	on	on	on	on	on	on	on	on	on	on	on	on	on	on	on	on	on	on	on	on	on	on	on	on	on	on	on	on	on	on	on

*NOTE--This treatment would not be successful where several pounds of seed are treated at a time, as the boiling temperature would not reach the center in the time allotted.

k---Killed
g---Good
w---Weak
p---Penicillium
r---Rhizopus
b---Bacteria
s---Sterigmatocystis

From Table VII it is seen that of all the temperatures tried, placing the seeds in boiling water for one or two seconds, seems to offer only little promise of success.

Placing the seeds in water of 90° C. for one minute insures a somewhat higher per cent of germination. However, the growth of seedlings seems to be weaker than either checks or those boiled for one or two seconds.

The series treated with water at temperatures of 80° C., 70° C., and 60° C. did not meet our expectations of success. However, these results are not final, as they simply open up a line of investigation for the future.

Effect of the Sulphuric Acid Treatment on Seed. Historical. Rostrup¹³³ was one of the first investigators to use sulphuric acid on hard seed in order to hasten germination. Todaro¹³⁴, while working independently, found that concentrated sulphuric acid of a density of 1.84 acted upon hard seeds of many leguminous plants, rendering them capable of prompt germination. Thornber¹³⁵, too, found that when certain seeds are treated with sulphuric acid, their germination was hastened. Schneider-Orelli¹³⁶ also found the sulphuric acid treatment of value in hastening the germination. Bolley¹³⁷ also found sulphuric acid to benefit the germination of seeds. In 1912 Love and Leighty¹³⁸ also found the same general beneficial results on germination of seeds treated with sulphuric acid.

My object in treating sweet pea seeds with sulphuric acid was to find out its effect on germination, and as a preventive means in destroying all possible adhering spores of pathogenic organisms. The method was to place the seeds to be treated in glass receptacles and then to cover the seeds with pure sulphuric acid. The time of treatment was five minutes, fifteen minutes, one-half hour, one hour and one and a half hours. After the treatment the acid was poured off and the glass receptacle was put under running tap water for five minutes and then rinsed three times in sterilized water. After that the seeds were placed on moist filter paper in petri dishes, and the latter were put in the incubator for ten days. A series of untreated seeds were also run as checks.

The results obtained from the seeds treated five minutes, fifteen minutes and one-half hour were practically the same, i. e., in each case the percentage of germination was much higher in the treated

seeds than in the checks. In the former the percentage of germination ranged from 95% to 100%, while in the later it ranged from 60% to 85%. The seeds treated in the acid for one hour showed 50% injury, and the 1½ hour treatment gave only 2% germination.

In order to test the effect of sulphuric acid on the fungus flora of the seeds, 10 cc of the acid was put in test tubes, and the latter were inoculated heavily with spores of *Glomerella rufomaculans* and allowed to stand for five minutes, fifteen minutes, half-hour and one hour. Transfers of the treated spores were made by means of a loop into melted tubes of agar. These were well shaken and poured into petri dishes, cooled, and placed in an incubator. Check cultures were also run by using untreated spores transferred directly into agar. In three days the check plates all showed a vigorous growth of a pure culture of the fungus where none of the series of the treated spores showed signs of germination even after eight days. This proves, then, that sulphuric acid can be used with advantage in treating seed both to increase the per cent of germination and also to kill all spores which adheres to the seed coat.

Formaldehyde Treatment of Seed. The method employed here was the same as for the sulphuric acid. The strength used was 5%, and the time of treatment was five minutes, fifteen minutes, half hour, one hour and one and a half hours. It was found that the one and a half hour treatment seemed to have reduced the percentage of germination, whereas, all the other treatments did not affect in any way the germination.

Where there was no injury apparent, the formaldehyde treatment did not seem to help the germination of the seeds as did the sulphuric acid. However, it no doubt helps to kill the adhering fungus spores of the seed coat. This latter advantage makes the formaldehyde treatment a valuable preventive means.

TREATMENT OF SOILS WITH CHEMICALS

The object of this treatment was to determine: 1, the effect of the treatment on the growth of the plant and its resistance to disease; 2, the effect on the soil flora; and 3, the effect on the nitrogen content and ammonification. The method employed was to sow 50 seeds in a pot (18 pots in all); the soil employed was unsterilized light garden loam.

The chemicals and the strengths used are indicated in Table VIII. After the seed had germinated and the plant had attained three inches in height, I began to water them twice a week with the respective chemicals. The checks were treated with distilled water. The experiment was run for two seasons, in each case up to the flowering time of the host.

The results of the first season are not indicated in Table VIII because with one exception as stated below there was no apparent difference between the treated and the check plants. During the first season, the treatment did not affect in any way the fungus or bacterial flora of the soil. Both treated and check plants, before the close of the first season, were inoculated with spores of *Glomerella rufomaculans*. But both lots gave about the same percentage of infection. This clearly indicated the wonderful power of the soil to absorb mineral poisons and to fix them in such a way as to make them harmless to plant growth. Ordinarily, neither plant, fungi, nor bacteria could grow in a solution of 1-1000 copper sulphate, for instance. However, when this same solution is applied to the growing plants through the soil, the latter fixes it so that the plants continue their growth and reach maturity as they would if the copper sulfate were not there. The same holds true for the soil flora.

The only injury apparent to the plants during the first season's trial was on the series watered with $1/100 \text{ MnSO}_4$. Altho growing fairly well, these plants were seen to lose their chlorophyll at an early date. These plants died just before blossoming, and at that stage they were white with no trace of chlorophyll.

The results obtained from the second season's growth are tabulated in Table VIII.

In order to determine the effect of the different chemical treatments on the soil flora, the method employed for isolating the organisms was the same as that recommended by Prof. T. F. Manns¹³⁹. Plates containing $1/1000$ and $1/10,000$ of a gram of soil was made. Two kinds of media were used for this purpose, the composition of which is given in Table VIII. The media marked XXI is purely synthetic and is of value in bringing out the azotifiers. It is also valuable in bringing out the bacterial flora of a soil and in keeping down the saprophytic fungi. Medium II on the other hand, is more likely to bring out the fungus growths and to keep in check the bacterial




TABLE VIII

Effect of treatment on plant growth during second season			Effect of treatment on soil flora			Effect of treatment on nitrogen content and ammonification x			
Chemical and dilution used	Nature of root growth	Nature of stem growth	Injury	Average bacteria per gram of soil	Average fungi per gram of soil	Total nitrogen dry basis	Per cent moisture	Quantity of N ₂ NH ₃ formed in 6 days	Quantity of N ₂ NH ₃ formed in 8 days
Cu SO ₄ 1-1000	E	C	70% of seedlings killed	Medium XXI* 1,140,000	Medium III† 390,000	0.2301%	14.25%	1.99cc	65.8cc
“ 1-2000	B	B	100% killed	3,130,000	160,000	0.2215%	14.43%	16.7 “	39.9“
“ 1-3000	B-E	A	100% killed	2,640,000	80,000	0.2354%	12.48%	13.8 “	31. “
Fe SO ₄ 1-1000	AA	A	—	1,550,000	30,000	0.1495%	15.75%	15.3 “	65.7“
“ 1-2000	AA	A	—	1,410,000	20,000	0.1444%	13.45%	25. “	58.6“
Kmn O ₄ 1-1000	AA	B	—	2,250,000	20,000	0.1432%	14.33%	16.1 “	61.3“
“ 1-2000	B	AA	—	1,430,000	20,000	0.1924%	12.2 %	20.6 “	65.5“
“ 1-3000	AA	B	—	890,000	20,000	0.1846%	13.2 %	4.8 “	35.6“
Hg Cl 1-2000	D-E	D	100% killed	3,640,000	30,000	0.1695%	19.5 %	16.4 “	55.9“
“ 1-4000	AA	B	—	3,690,000	20,000	0.1963%	19 %	8.9 “	59.4“
Zn SO ₄ 1-100	toproots	C	100% killed	2,420,000	90,000	0.1813%	12.2 %	1.6 “	44.5“
“ 1-500	D-E	B	—	3,580,000	20,000	0.1817%	20.25%	6.2 “	36.9“
“ 1-1000	B	B	—	2,380,000	20,000	0.1912%	16 %	7.2 “	62.9“
Mn SO ₄ 1-100	E	B	100% killed	1,350,000	60,000	0.1750%	17.6 %	6.7 “	43.8“
“ 1-500	A	A	—	1,110,000	10,000	0.1803%	18.5 %	14.2 “	45. “
“ 1-1000	AA	A	—	1,890,000	20,000	0.1754%	—	—	—
Check, untreated soil	A	A	—	2,800,000	30,000	0.1984%	13.5 %	8.4 “	45.4“
Nutrient broth not inoculated check	—	—	—	—	—	—	—	2.2 “	2.4“

AA Stimulation in growth

A Good Growth

B Medium Growth

C Poor Growth

D Very poor growth

E Killed by treatment

Remarks: For stem and root growth, the check was taken as a standard of comparison.

x Note: Ammonification by Remy's method, using a peptone solution and a soil infusion. Not a reliable method.


* Medium XXI is made as follows:

Agar	15 grams
Mannite	10 “
Magnesium sulphate	0.2 “
Basic slag	4. “
Water (distilled)	1000 cc

The above was taken from Florence Mackeridge, Kings College, London 1912. This medium is good for isolation and growing of Azotobacter.

† Medium II is made as follows:

Glucose	20 grams
Peptone	10 “
Dipot. phosphate	0.25 “
Magnesium sulphate	0.25 “
Agar	15 “
Water	1000 cc



flora. The nitrogen was determined* according to the Kjeldahl (modified) method₁₄₀.

Ammonification was determined by inoculating 100 cc of nutrient broth containing one gram of peptone with 1 cc of an infusion made by shaking 10 grams of soil in 100 cc of water.

From Table VIII we see that the results obtained vary considerably, especially during the first six days, which more nearly represents the ammonifying power. Under copper sulphate for instance, 1/1000 killed 70% of the seedlings and the growth of the latter was quite stunted. The average number of bacteria per gram of soil was also less when compared to the checks, whereas there was an actual increase in the nitrifiers, which resulted in an increase of total nitrogen and ammonia. On the other hand, copper sulphate 1/3000 produced stimulation in plant growth; there did not seem to be an appreciable decrease in the general bacterial flora, but there was a resultant decrease of ammonifiers. Again, under Hg Cl₂ 1/2000, the result was a killing of all the seedlings, but an increase in the bacterial flora as well as nitrification and ammonification. Hg Cl₂ 1/4000 gave stimulation in plant growth, soil flora, and in ammonification.

Table VIII is extremely interesting, as it opens up so many new phases in soil biology, soil bacteriology, and in plant pathology.

STUDIES OF THE FUNGICIDAL VALUE OF SOME CHEMICAL POISONS

Those who are actively engaged in the study of methods of control in plant disease, will readily realize how uncertain it is to depend upon field methods alone. In order to test out the value of a fungicide the only method used consisted in spraying the particular plant to be treated, with that fungicide, and of noting results. With such a method one works in the dark. Moreover, a great deal of time is lost, because each trial, or each modification of a trial means one season of growth. In 1910 Reddick and Wallace₁₄₁ worked out a quick method of determining the efficiency of a certain spray material. In brief, the method is as follows:

The fungicide to be tested is sprayed on a clean slide with an atomizer. It is then allowed to dry in order to permit any chemical

*Mr. Paul Emerson under the direction of Prof. Manns carried out this phase of the work.

changes to occur that are likely to be induced by exposure to atmospheric conditions. It is supposed that the same chemical changes which take place on the leaf previous to infection also take place on the slide in the laboratory before the spores are placed to germinate. This more nearly duplicates natural conditions, and the results at time of germination practically indicate the relative fungicidal value of the chemical tested. When thoroughly dry, each slide is placed in a petri dish containing enough water to keep the air well saturated. Spores of the particular fungus which we wish to control are mixed in water, and drops of this water are placed, with a dropper, on the treated slide. The petri dish is now covered and 24 hours are allowed for the spores to germinate. The inhibition of germination of all the spores in the drops placed on the slide indicates the value of the fungicide. While the method above described is excellent, it nevertheless has its drawbacks. We know that in practice we do not spray our plants every day, but this is usually done every three or four weeks. The method employed by Reddick and Wallace cannot account for the fungicidal value of a certain spray material during the two or three weeks from the time of its first application. In my studies I have supplemented this deficiency by the following method. While adopting the method used by Reddick and Wallace of applying the fungicide and of drying the slide, I have worked out the following modifications. Instead of spraying only a few slides and of inoculating them at once, I have sprayed some fifty slides at one time with each chemical. These slides were all dried and divided into different lots. Lot 1 was used for germination at once, lot 2 used, say, after an interval of one week, etc. This method allows the time element to have its full effect on the fungicide concerned, and each germination test definitely indicated whether that fungicide deteriorates rapidly or not. The fungicide which preserves its germicidal value the longest is probably of the greater economical value. The fungicides tested and the strengths used are indicated in Table IX.

It will be seen from the above table, that copper sulfate at the strength of 1/100 up to 1/500, and potassium permanganate 1/2% up to 3% completely inhibited the germination of the spores of *Glomerella rufomaculans* for nearly two months. On the other hand, lime sulphur 1/10 up to 1/50 gave most unsatisfactory results from the very first test, and its fungicidal value rapidly weakened with age. This was still more evident with the "Sulphurated Potassium" also

TABLE IX

A		B		C		D	
Strength of poison used	Date of inoculating the slides	Per cent of spore germination	Strength of poison used	Date of inoculating the slides	Per cent of spore germination	Strength of poison used	Date of inoculating the slides
Cu SO ₄ , 1-100	3-7-12	0	Potassium permanganate 1/2%	3-7-12	0	Lime Sulphur 1-10	3-7-12
" " 1-200	" "	0	" " 1%	" "	0	" " 1-20	" "
" " 1-300	" "	0	" " 2%	" "	0	" " 1-30	" "
" " 1-400	" "	0	" " 3%	" "	0	" " 1-40	" "
" " 1-500	" "	0	Check	" "	92%	Check	" "
Check	" "	100%	Pot. per. 1/2%	4-18-12	0	L. Stul. 1-10	3-13-12
Cu SO ₄ , 1-100	3-13-12	0	" " 1%	" "	0	" " 1-20	" "
" " 1-200	" "	0	" " 2%	" "	0	" " 1-30	" "
" " 1-300	" "	0	" " 3%	" "	0	" " 1-40	" "
" " 1-400	" "	0	Check	" "	100%	Check	" "
" " 1-500	" "	0	Pot. per. 1/2%	3-26-12	0	L. Stul. 1-10	3-26-12
Check	" "	100%	" " 1%	" "	0	" " 1-20	" "
Cu SO ₄ , 1-100	3-26-12	0	" " 2%	" "	0	" " 1-30	" "
" " 1-200	" "	0	" " 3%	" "	0	" " 1-40	" "
" " 1-300	" "	0	Check	" "	100%	Check	" "
" " 1-400	" "	0	Pot. per. 1/2%	3-13-12	0	L. Stul. 1-10	4-18-12
" " 1-500	" "	0	" " 1%	" "	0	" " 1-20	" "
Check	" "	100%	" " 2%	" "	0	" " 1-30	" "
Cu SO ₄ , 1-100	" "	0	" " 3%	" "	0	" " 1-40	" "
" " 1-200	" "	0	Check	" "	82%	Check	" "
" " 1-300	" "	0	Check	" "	0	" " 1-20	" "
" " 1-400	" "	0	" "	" "	0	" " 1-30	" "
" " 1-500	" "	0	" "	" "	0	" " 1-40	" "
Check	" "	100%	" "	" "	0	Check	" "
Cu SO ₄ , 1-100	" "	0	" "	" "	0	" "	" "
" " 1-200	" "	0	" "	" "	0	" "	" "
" " 1-300	" "	0	" "	" "	0	" "	" "
" " 1-400	" "	0	" "	" "	0	" "	" "
" " 1-500	" "	0	" "	" "	0	" "	" "
Check	" "	100%	" "	" "	0	" "	" "

known as "Liver of Sulphur." Lime sulphur is a valuable fungicide in controlling apple scab (*Venturia inequalis*) and sulphurated potassium is a valuable fungicide in controlling mildew (*Oidium* sp.), but evidently these two fungicides are of no value in the control of the sweet pea anthracnose. On the other hand, copper sulphate and potassium permanganate both prove very toxic to the spores of the anthracnose. The next step therefore, was to test all the strengths used of these two fungicides by spraying them on sweet pea plants in field, and by watching to see if these poisons produced burning of the leaves and stems. Copper sulphate at the strengths of 1/100, 1/200, 1/300, 1/400 and 1/500 all burned the leaves of the sweet pea. Hence while the above strengths possess valuable germicidal properties their use upon the sweet pea is prohibitive. This means that weaker strengths are to be tried until the proper limit is reached, i. e. the limit which does not decrease the germicidal value of the copper sulfate and which does not produce injury to the plant. This we intend to carry on further in the future. Potassium permanganate $\frac{1}{2}\%$ up to 3% does not produce any injury to the plant whatsoever. As a matter of economy, therefore, $\frac{1}{2}\%$ of potassium permanganate proves a valuable fungicide in the control of the sweet pea anthracnose. It should be applied to the plant not oftener than it is washed off by rain. The method can be used in testing out an endless number of chemicals and there is no doubt that a good many will prove valuable additions to our list of fungicides.

SOIL TREATMENT IN THE GREENHOUSE

Growers who are troubled with *Rhizoctonia* or *Fusarium* in their greenhouse beds, will find the following directions valuable¹⁴².

STEAM STERILIZATION

"The preventive method which promises best results to those who have the convenience for applying it is that of sterilization of the seed beds by steam.

"In addition to the killing of the fungus, this method, in common with surface firing, to be described later, has several advantages over formalin treatments. The weed seeds in the soil are very largely killed, and this alone, according to the testimony of the farmers who have used sterilization, pays for the cost of treatment, as the beds do not have to be weeded and thus a large amount of hand labor is obviated. The physical texture of the soil is altered by the heat and

made more suitable to root development and, moreover considerable plant food is made directly available to the seedlings. Furthermore, the heating of the soil just before sowing in the spring has an appreciable effect in starting the seedlings off quickly.

“With the elimination of the fungus it is possible to employ those methods forcing the plants by extra fertilization, increased watering, and higher temperature which would otherwise be unsafe as favoring the development of the root-rot fungus.

“*Ordinary greenhouse method*—The method of sterilization to be used will depend to some extent on the size, the location, and the permanency of beds and the cost of application.

“The method in general use for the sterilization of soil in greenhouse benches might advantageously be employed in beds that are to be used year after year without change of location, as the equipment would be more or less permanent. This consists in placing one foot below the surface of the soil a system of $1\frac{1}{2}$ -inch pipes which are perforated with $\frac{1}{4}$ -inch holes on their under side at intervals of 6 inches throughout their entire length. The pipes should run lengthwise of the bed, 18 inches apart, and be connected with a steam boiler capable of producing 80 to 100 pounds pressure. Before treatment the soil should be thoroughly spaded up and pulverized to permit ready access of the steam to all parts, and all fertilizers should be applied at this time.

“The bed to be treated should be covered with several thicknesses of old burlap or blankets to confine the heat to the soil. The steam should be applied at a pressure of 80 to 100 pounds, as at a high pressure it is much drier and the soil is not wet as much as when low-pressure steam is used. A treatment of from one to two hours is usually sufficient to thoroughly sterilize the soil to a depth of 18 inches. A few potatoes laid in the surface will indicate the thoroughness of the treatment by the degree to which they are cooked. The blankets might advantageously be left on for some time to make the treatment more thorough.

“While this method offers some advantages for seed beds of limited area, in that the pipes may be left in the ground and used year after year with little extra labor and may also be used for subirrigation, the initial cost of installation, especially on large seed-bed areas, may be prohibitive.

“Inverted-pan method. The method which has given the best results in practice, and which because of its simplicity and small cost recommends itself for use on large or small areas, is the invention of Mr. A. D. Shamel, of the Bureau of Plant Industry, and was devised by him to sterilize nematode-infested soils in Florida. The apparatus consists of a galvanized iron pan, 6 by 10 feet and 6 inches deep, which is inverted over the soil to be sterilized and the steam admitted under pressure. The pan is supplied with steam hose connections, has sharp edges, which are forced into the soil on all sides to prevent the escape of steam, and is fitted with handles for moving it from place to place, the weight of the entire pan being not over 400 pounds.

“The soil is prepared as in the greenhouse method, a few potatoes being buried at a depth of a foot to gauge the degree of heat attained. A soil thermometer may also be used if desired. The steam should be kept at as high a pressure as possible, 80 to 100 pounds being best, and the treatment should continue for one to two hours, depending on the pressure maintained. In experiments conducted in the spring of 1907, one hour’s steaming at 80° C. under 100 pounds pressure gave best results in killing both the fungus and the weed seeds. When one section of the bed is treated the pan is lifted and carried to an unsterilized portion and the operation repeated until the entire bed is steamed.

FORMALDEHYDE STERILIZATION

“The use of a formalin solution for the sterilization of greenhouse soil against *Rhizoctonia* has been in vogue for some time with excellent results, and furnishes a very simple means of combating the root-rot. The method is as follows: The beds are thoroughly prepared the same way as for the other methods of sterilization described and are then drenched with a formalin solution composed of 1 part of commercial formalin to 150 to 200 parts of water, three-fourths to 1 gallon of this solution being used to the square foot of bed space. The solution should be put on with a watering pot with a hose and distributed as evenly as possible over the bed, so as to thoroughly wet the soil to the depth of a foot. It will in most cases be necessary to put this solution on in two or three applications, as the soil will not take in this quantity of water immediately. The beds should then be covered with heavy burlap or a tarpaulin to keep in the fumes for a day or so, and then aired for a week before sowing the seed.

“Spring applications of formalin are open to the following objections: The addition of such a large quantity of water to the soil keeps it wet and cold for some time longer than would naturally be the case, thus delaying germination as well as subsequent growth; the necessity of airing the beds to remove the formalin fumes and to allow the soil to dry out also causes delay in seeding. To obviate this difficulty the beds should be treated in the fall, before freezing weather sets in. In this case a stronger solution, 1 to 100, may well be used, as there will of course be no danger then of injuring the seedlings.”

SUMMARY

1. It has been shown that the sweet pea is subject to a number of diseases.

2. The following classes of diseases have been investigated: I. Fungous; II. Bacterial; III. Physiological; IV. Animal or Insect Pests.

3. Contrary to the statements of Masee and Chittenden, *Thielavia basicola* does not produce the “streak,” but produces only a root rot of the sweet pea.

4. The pathogenic nature of *Corticium vagum* B. & C. has been established.

5. *Chaetomium spirochaete* has been shown for the first time to be a plant pathogen, and especially to produce a root rot of the sweet pea.

6. A new *Fusarium* root disease has been described, and the name *Fusarium lathyri* Taubenhau has been given to the fungus, and its pathogenicity established.

7. Of the Animal parasites the eel worm (*Heterodera radicolica*) has been shown to produce a root gall disease of the sweet pea. The disease is carefully described. The eel worm has also been shown to open the way to the attacks of several fungous diseases.

8. *Sclerotinia libertiana* has been shown for the first time to produce a collar rot as well as a stem disease of the sweet pea.

9. Studies on the mildew of the sweet pea have shown the disease to be very prevalent under greenhouse conditions as well as out of doors. The cause of the mildew is a species of *Oidium*. Observations up to date have failed to reveal the perfect stage of the fungus.

10. Extended studies and cross inoculations have definitely proven that the anthracnose disease of the sweet pea and the bitter rot of the apple are caused by the same fungus *Glomerella rufomaculans*.

11. It has been also proven that the following pathogens, namely, *Gloeosporium gallarum* from oak gall, *Gloeosporium diospyri* from persimmon fruit, *Gloeosporium officinale* from the Sassafras, *Colletotrichum nigrum* from the pepper plant, *Colletotrichum phomoides* from the tomato, appear to be identical and the same as *Glomerella rufomaculans*, since they can all produce the anthracnose disease of the sweet pea and the bitter rot of the apple. Cross inoculations would no doubt reduce the great number of our so-called different species of *Gloeosporium*s.

12. The mosaic has been shown for the first time to produce a disease on the sweet pea. The pathogenicity and the infectious nature of the disease have been clearly demonstrated. Exceptions are taken with Woods that the disease is of a physiological nature but that it is induced by either bacteria or protozoa which neither our microscope nor our present method of staining are able to detect. Insects and especially the green aphids seem to be the main carriers and distributors of the disease.

13. Manns and Taubenhaus have definitely proven that the "streak" disease is caused by *Bacillus lathyri* M. & T. and not by *Thielavia basicola* as previously believed by Masee and Chittenden.

14. Bud Drop in one form is here shown to be induced by a high nitrogen supply not properly balanced by phosphoric acid and potash; by addition of the latter the trouble is quickly corrected. Arrested development may be due to overtreatment of soil with wood ashes the treatment being too caustic. Such an error may be corrected by the addition of acid phosphate.

15. Under methods of control it is shown that no one variety is immune to the anthracnose but there are certain individuals in each variety which are more or less immune to the disease.

16. Boiling the seeds for one or two seconds destroys the spores of parasitic fungi, but commercially the treatment is not applicable on large quantities of seed at a time.

17. Soaking the seeds in sulphuric acid for five minutes, fifteen minutes and one-half hour increases the per cent of germination and at the same time kills all the spores which adhere to the seed coat.

18. Soaking the seeds in a 5% formaldehyde solution from five minutes to one hour does not increase nor decrease the percent of germination but helps to kill the spores which adhere to the seed coat.

19. Watering soils with chemical poisons does not increase the resistance of the plants which are grown on that soil. The latter adsorbs and fixes some of these poisons so as to make them harmless to plant growth.

20. A new method has been devised in determining the length of time in which any fungicide can remain efficient in controlling plant diseases when sprayed on the plant to be treated.

REFERENCES

1. Condensed from A. C. Beal. Sweet pea studies III. Cornell Agr. Expt. Sta. Bul. 320: 661-713. 1912.
2. Bailey, L. H., Cyclopedia of American Horticulture, p. 1752. 1902.
3. Cuthbertson, Michael, Sweet pea cupid, Gard. Chron., Vol. XX, third series: 404, 1896.
4. Masee, George. Fungoid diseases of the sweet pea. Sweet Pea Annual: 18-22, 2 figs. 1906 (London).
5. Weston, T. A., The "Streak". The Gardener, 1906.
Weston, T. A. The streak disease of sweet peas. The Gardener: 585, 1907.
6. Anonymous. Sweet pea diseases. The Fruit Grower, Fruitier, Florist and Market Gardener: 295, 1907.
7. Masee, George. Sweet pea streak. Sweet Pea Annual: 22, 1909.
8. Masee, George. A disease of sweet peas, asters and other plants. Roy. Bot. Garden Kew Bul. of Misc. Information, No. 1: 44-52, 1 plate, 9 figs. 1912
9. Chittenden, F. J. Diseases of sweet peas, Sweet Pea Annual: 14-24, 1912 (London).
10. Dyke, W. Sweet pea streak. Amateur Gardening: 593, 1912.
11. Sheldon, J. L. Concerning the identity of the fungi causing an anthracnose of the sweet pea and the bitter rot of the apple. Science N. S. 22: 51-52. 1905.
12. Taubenhaus, J. J. A Study of some Gloeosporiums and their relation to a sweet pea disease. Phytopathology 1: 196-202, 1911.
Taubenhaus, J. J. A further study of some gloeosporiums and their relation of a sweet pea disease. Phytopathology 2: 153-160, 1912.
13. Taubenhaus, J. J., and Manns, T. F. Diseases of the sweet pea. Gard. Chron. Vol. LIV: 21-26. 1913 (London).
14. Berkeley, M. J., and Broome, C. E. Notices on British fungi. Ann. and Mag. Nat. Hist. s. 2, Vol. 5: 461, 1850. London.
15. Zopf, W. Thielavia, genus novum, Perisporiacearum. Verhandl. Bot. Ver. Prov. Brandenburg j. 18. Sitzungsber 30 juni, 1876, pp. 101-105. Berlin, 1876.
16. Sorokin, N. Ueber Helminthosporium fragile sp. n. Hedwigia. Bd. 15: 113-114, 1876.
17. Saccardo, P. O. Clasterosporium fragile (Sorock.) Sacc. Sylloge Fungorum, 4: 386, 1886.
18. Sorauer, P. Handbuch der Pflanzenkrankheiten (Auf. 2) 1886. Berlin.
19. Loc. cit. p. 9.
20. Loc. cit. p. 8.
21. Loc. cit. p. 9.

21. Peglion, V.—Atti R. Accad. Lincei, an 294, s. 5, Rend. Cl. Sci. Fis., vol. 6, semestre 2, pp. 52-56. Roma, 1897.
22. Aderhold R. Impversuche mit *Thielavia basicola* zopf. Arb. Biol. Abt. Land. Forthw. Kaiserl. Gesundh. 4: 463-465. 1905.
23. Clinton, G. P. Root rot of tobacco. Conn. Agr. Expt. Sta. Rept. 1907.
24. Gilbert, W. W. The root rot of tobacco caused by *Thielavia basicola*. Bur. Plant Indust. Bul. 158, 1909, Washington, D. C.
25. Manns, T. F. Studies in diseases of cereals and grasses. Ohio Agr. Expt. Sta. Bul. 203. 1909.
26. Duhamel, H. L. Explication physique d'une maladie qui fait perir plusieurs plantes dans le Gatinois et particulierement le Safran. Mem. Acad. Sciences, Paris, 1728.
27. Fougereux de Bandaroy, Hist. de l'Acad. Roy. d. Sc. Annee 1782, p. 89, 1785, Paris.
28. Builliard, P. Histoire des champignons de la France T. I. p. 81, 1791, Paris.
29. Persoon, C. H. Synopsis methodica fungorum Gottingae p. 119, 1801.
30. De Candolle, A. P. Memoire sur les Rhizoctones. Mem. de Museum d' Histoire naturelle T. II, p. 209, 1815, Paris.
31. Nees, von Esenbeck. Th. F. L. System der Pilze Schamme p. 148, 1817.
32. Duby, H. E. Botanicum gallicum. Ed. II, Paris, p. 867, 1830.
33. Leveille, J. H. Mem. sur le genre *Sclerotium*. Ann. d. sc. nat. bot. T. XX, 1843, Paris.
34. Tulasne, L. R. and C. Fungi hypogaei p. 1851, Paris.
35. Kuhn, J. Krankheiten des Kulturgevachse, 1858.
36. Erikson, J. Einige Studies uber den Wurzeltoter (*Rhizoctonia violacea*) des Mohre mit besonderer Rucksicht auf seine Verbreitungs. fahigkeit Centrallblatt f. Bakt. Abh. II, Vol. X: 721 & 766, 1903.
37. Gussow, Th. Jour. Roy. Agr. Soc. England. p. 175, 1905.
38. Shaw, F. J. F. The morphology and parasitism of *Rhizoctonia*. Mem. Dept. Agr. in India. Bot. Vol. IV, pp. 115-153, 2 plates. 1912, India.
39. Pammel, L. H. Bul. 15, Iowa Agr. Exp. Sta. 1891.
40. Atkinson, G. F. Some diseases of cotton, Bul. 41. Ala. Agr. Expt. Sta. 30-39, 1892.
41. Stone, G. E., and Smith, R. E. The rotting of greenhouse lettuce. Mass. Agr. Expt. Sta. Bul. 186, 1901.
42. Duggar, B. M., and Stewart, F. L. The Sterile fungus *Rhizoctonia*, Cornell Agr. Expt. Sta. Bul. 186, 1901.
43. Rolfs, F. M. Potato failures. A second rept., Colo. Agr. Expt. Sta. Bul. 91, 1904.
44. Stevens, F. L., and Hall, J. C. Hypochnose of pomaceous fruits. Ann. Mycologici, Vol. VII, 1909. Diseases of economic plants, 1910.
45. Fuckel, L. Symbolae mycologicae. Wiesbaden, p. 142, 1869.
46. Prunet, A. Sur le *Rhizoctonia* de la lucerne. Compt. Rend. T. C. XVII 1893, Paris.
47. Hartig, R. Untersuch aus d. forstbotan. Institut. zu Munchen, 1888.

48. Frank, B. Ein neuer Rebenbeschädiger in Rhenihessen. Zeitsch. f. d. Landw. Vereine d. Gossh. Hessev. 1897. No. 19, p. 167.
49. Loc. cit. p. 19
50. Loc. cit. p. 19.
51. Reinke J. und Berthold G. Die Zersetzung Der Kartoffel Durch Pilze. Berlin, 1879. pp. 100, tables XI.
52. Loc. cit. p. 26
53. Tulasne. Selecta fungorum carpologia tom. III. 1865.
54. Hansen, Ch. Dr. Jour. Roy. Micr. Soc. 321. 1883.
55. Smith, Worthington G. Diseases of field and garden crops, 1884.
56. Smith, E. F. The fungus infestation of agricultural soils in the United States. Proc. Amer. Assoc. Adv. Sci. 48: 303, 1889.
57. Kellerman, W. S., and Swingle, W. T. Loose smuts of cereals. Second Ann. Report, Kansas State. Agr. Expt. Sta. 213-238. 1889.
58. Frank, Jahb. d. deutsch landw. Ges. 1892.
59. Atkinson, G. Some diseases of cotton. Ala. Agr. Experiment Sta. Bul. 41, 1892.
60. Rostrup. Landboskrifter, Bd. V, 1893.
61. Smith, E. F. Wilt disease of cotton, watermelon and cowpeas. Bu. Plant Indus. U. S. Dept. Agr. Bul. 17, 1899.
62. Woods, A. F. The aster disease. Gardening 7: 277, 1899.
63. Manguin, Louis. Sur le parasitisme *Fusarium roseum* et de especes affinie.
64. Prillieux et Delcroix. Sur la maladie des oeilletts produits par le *Fusarium dianthi* P. & Del.
65. Bolley, H. L. Flax wilt and flax sick soil. N. D. Sta. Bul. 50, 27-60, 1901.
66. Sorauer, P. Pflanzenkrankh. 11: 217-228, 1901.
67. Pammel, The grasses of Iowa. 1901.
68. Smith, R. E. Growing china asters. Mass. Sta. Bul. 79, 1902.
69. Loc. cit. p. 28.
70. Hennings. P. Ztschr. Pflanzenkrankh. 12: 14-16, 1902.
71. Hall, Van C. Ber. d. Deutsch. bot. Ges. 1903, p. 2.
72. Smith, E. F., and Swingle. The dry rot of potatoes. Bur. of Plant Industry, U. S. Dept. Agr. Bul. 55, 1904.
73. Osterwalder, A., Dr. Über eine buscher unbekannte Art. der kernobstfaule verursacht durch *Fusarium putrefaciens* Centbl. Bakt. 2 Abt. 207-213, 1904.
74. Owen, V. Über ein *Fusarium* Krankheit der Tomaten. Landw. zährb. p. 489, 1905.
75. Appel, O. und Schikarra, G. Arb. K. Biol. Inst. Land. U. Forstw. 5: 155-188, 1906.
76. Heald, F. D. Bud rot of carnations Abs. in Sci. n. ser 23: 620, 1906.
77. Hedgecock, G. G. Studies upon some chromogenic fungi which discolor wood. Mo. Bot. Garden Rept. 17. III. 1906.
78. Chiffot, J. A. A study of disease of pelargonium Journ. Soc. Nat. Hort. France 9: 348-355, 1907.

79. Burrill, T. J., and Barrett, J. T. Ear rots of corn. Illinois Sta. bul. 133, 1909.
80. Wolf, F. A. A *Fusarium* disease of the pansy. *Mycologia* 2: 19-22, 1910.
81. Smith, E. F. A Cuban banana disease. Abs. in Science N. ser 31: 754-755, 1910.
82. Cook, M. T. The double blossom. Abs. in Sci. N. ser 751, 1910.
83. Bubak, J. F., and Kosaroff, P. Some interesting plant diseases from Bulgaria, *Centbl. Bakt.* 2 abt. 31: 495-502, 1911.
84. Gifford, C. M. The damping off of coniferous seedlings. *Vt. Sta. Bul.* 156: 143-171, 1912.
85. Wollenweber, H. W. Studies on the *Fusarium* problem. *Phytopathology* 3: 24-50, 1913.
86. Loc. cit. p. 29.
87. Marcinowski, Kati. Parasitisch und semi parasitsch an Pflanzen lebende nematoden Arbeiten aus der K. Biol. Anst. fur Land und Forstwirtschaft, Berlin 7: 1-192. 1909.
88. Bessey, E. A. Root knot and its control. U. S. Dept. Agr., Bureau Pl. Industry., Bul. 217: 7-82, 1911.
89. Loc. cit. p. 32.
90. Chittenden, F. J. Diseases of the sweet pea. *Sweet Pea Annual*, 14-24, 1912. England.
91. Coemans, M. E. Recherches sur la genese et les metamorphoses de la *Peziza sclerotiorum* Lib. *Bul. l'acad. roy. des Sciences de Belgique* II, 9: 61, 1860.
92. De Bary, A. Uber einige Sclerotinien und Sclerotinien Krankheiten *Bot. Zeitz* 44: 378, 1886.
93. Cohn, F. *Illustr. landw. Ztg.* 1887.
94. Smith, W. G. *Gard. Chron.* III, 8: 324, 1890.
95. Behrens, J. Uber das auftreten des Hanf Krebses in Elsass. *Zeitschr der Pflanzen Krank* 1: 208. 1891.
96. Humphrey, J. E. The rotting of lettuce. *Rept. Mass. State Expt. Sta.* 8: 219. 1892.
- Humphrey, J. E. Disease of the cucumber plant. A sclerotium disease. *Rept. Mass. Txpt. Sta.* 10: 212, 1893.
97. Prillieux, M. M. et Delacroix, *Maladie des branches des muriers de la Turquie d' Europe.* *Compt. Rend Acad. Sci., Paris.* Vol. 124: 1168-1176, 1897.
98. Smith, Ralph E. *Botrytis and Sclerotinia. Their relation to certain plant diseases and to each other.* *Bot. Gaz.* 29: 369-406, 1900.
99. Hedgecock, G. G. A disease of cauliflower and cabbage. *Mo. Bot. Gard. Ann. Rept.* 16: 149-151, 1905.
100. Parisot, F. *Maladie des Topinambours, Jour. Agr. Prot. N. Ser.* 10: 369-371, 1905.
101. Appel, Otto, and Bruch, F. W. *Sclerotinia libertiana* Fekl. als Schadiger von Wurzel Fruchten *Arb. K. Biol. anst. Land u. Forstw.* 5: 189-263, 1906.
102. Masseron, P. Une nouvelle maladie des pois cultives. *Semaine Agricole (Paris)* 26. 270, 1907.

103. Westerdijk. Zoha. Untersuchungen über *Sclerotinia libertiana* Fekl. als Pflanzenparasit mended. Phyto. Lab. Willie Commelin Scholten. pp. 5-26. 1911.
104. Taubenhaus, J. J. Present knowledge of sweet pea diseases. Florist exchange 34: 108-110, 1912.
105. Loc. cit. p.
106. Masee, G. Fungoid diseases of the sweet pea: Sweet Pea annual, 20-21, 1906.
107. Stewart, F. C. Notes on New York plant diseases. New York (Geneva) Agr. Expt. Sta. Bul. 328: 394, 1910.
108. Sheldon, J. L. Concerning the identity of the fungi causing an anthracnose of the sweet pea and the bitter rot of the apple. Science, N. S. 22: 51-52. 1905.
109. Edgerton, C. W. Diseases of the fig and fruit. La. Agr. Exp. Sta. Bul. 126: 7. 1911.
110. Cook, M. T., and Taubenhaus, J. J. The relation of parasitic fungi to the contents of the cells of the host plants. Del. College Agr. Expt. Sta. Bul. 97: 3-52. 1912.
111. Mayer, Adolf. Landw. Versuchsstation, Vol. 32, 1886.
112. Land- und Forstwirtschaft, 1892. Cf. Berh. Z. Bot. Centralbl. Bd. 111, p. 266.
113. Comp. rendu hebdomadaire des Seances de l'Académie des Sciences. CXVIII-XII, p. 668, 1894.
114. Revue mycologique XIX, 73, p. 13, 1897.
115. Loc. cit. p. 53.
116. Woods, A. F. Observations on mosaic diseases of tobacco. U. S. Dept. of Agr. Bur. of Pl. Industry. Bul. No. 18, 1902.
117. Selby, A. D. Tobacco disease and tobacco breeding. Ohio Agr. Expt. Sta. Bul. 156, 1904.
118. Allard, H. A. The mosaic diseases of tobacco. Science N. ser. Vol. XXXVI: 875-876, 1912.
119. Loc. cit. p. 10.
Exchange, 1912.
120. Loc. cit. p. 53.
121. Loc. cit. p. 53.
122. Conn. Agr. Expt. Sta. Rept. 1898.
123. Loc. cit. p. 54.
124. Bulletin of the College of Agr., Imp. Univ. Tokyo, IV, No. 3, 1900.
125. Masee, George. Fungoid diseases of the sweet pea. Sweet Pea Annual, 18-22, 1906.
126. Weston, T. A. Gardener, 1906.
127. Masee, George. The Sweet Pea Annual, 1909, England.
128. Chittenden, F. J. The Sweet Pea Annual, 14-24, 1912.
- Chittenden, F. J. On some plant diseases new or little known in Britain. Jour. Royal Hort. Soc. 36: 545-550, 1912.

129. Masee, George. A disease of sweet peas, asters, and other plants (*Thielavia basicola* Zopf.) Bul. of Miscel. Information: 44-52, 1912. Kew Bot. Garden, London.
130. Dyke, Van. Gard. Chron. 51: 36; 52-53; 84-85, 1912. London.
131. Masee, George. Sweet Pea Annual, 1912.
132. Manns, T. F., and Taubenhaus, J. J. Streak a bacterial disease of the sweet pea and clovers. The Gard. Chron. (London), Apr. 5, 1913, pp. 215-216.
133. Rostrup, O. Report of the Danish seed control for 1896-97. pp. 37, 1898. Copenhagen (Review E. L. R. 10: 53-54.)
134. Todaro, F. Azione dell' Acido solforico concentrato su alcuni semi, 2 in particolare sopra i semi duri delle Leguminose. Staz. Sper. Agr. Ital. 34: 613-689, 1911 (Review E. S. R. 13: 754-755.)
135. Thornber, J. J. Arizona Agr. Expt. Station, Rept. pp. 489-493, 1904.
136. Schneider-Orelli, O. The resistance of medicago seed to high temperatures. Flora 100: 305-311, 1910 (Review E. S. R. 24: 231.)
137. Bolley, H. L. The agricultural value of hard seeds in alfalfa and clover seeds. Paper read before the Association of Official Seed Analysis, 1910.
138. Love, Harry H., and Leighty, Clyde E. Germination of seeds as affected by sulphuric acid treatment. Cornell Univ. Agr. Expt. Sta. Bul. 312: 295-336, 1912.
139. Manns, T. F. The Fungi of flax sick soil and flax seed (unpublished) 1903, and later methods by the same author.
140. Bul. 107, Bur. of Chem.
141. Reddick, D., and Wallace, E., Science, n. ser. 31: 789, 1910.
142. Quoted from Gilbert, W. W. The root rot of tobacco caused by *Thielavia basicola*. U. S. Dept. Agr., Bur. Pl. Industry., Bul. 158: p. 48, 1909.





LIBRARY OF CONGRESS



0 021 468 248 5